arXiv:1302.3744v2 [math.RT] 12 Mar 2013

LOCAL THETA LIFTING OF GENERALIZED WHITTAKER MODELS ASSOCIATED TO NILPOTENT ORBITS

RAUL GOMEZ AND CHEN-BO ZHU

Abstract. Let $(G, \tilde{G})$ be a reductive dual pair over a local field $\Bbbk$ of characteristic 0, and denote by V and $\tilde{V}$ the standard modules of G and $\tilde{G}$, respectively. Consider the set $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ of full rank elements in $\operatorname{Hom}(V, \tilde{V})$, and the nilpotent orbit correspondence $\mathcal{O} \subset \mathfrak{g}$ and $\Theta(\mathcal{O}) \subset \tilde{\mathfrak{g}}$ induced by elements of $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ via the moment maps. Let $(\pi, \mathscr{V})$ be a smooth irreducible representation of G. We show that there is a correspondence of the generalized Whittaker models of π of type $\mathcal{O}$ and of $\Theta(\pi)$ of type $\Theta(\mathcal{O})$, where $\Theta(\pi)$ is the full theta lift of π. When $(G, \tilde{G})$ is in the stable range with G the smaller member, every nilpotent orbit $\mathcal{O} \subset \mathfrak{g}$ is in the image of the moment map from $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$. In this case, and for $\Bbbk$ non-Archimedean, the result has been previously obtained by Mœglin in a different approach.

Contents

1. Introduction and main result 2
2. Classical groups as isometry groups of Hermitian modules 6
2.1. Hermitian D-modules 6
2.2. Isometry groups 7
3. Nilpotent orbits of classical groups 8
3.1. Nilpotent orbits and $\mathfrak{sl}_2$-triples 8
3.2. ϵ-Hermitian Young tableaux 9
3.3. Generalized Whittaker models associated to nilpotent orbits 11
4. A realization of generalized Whittaker models: $\Bbbk$ Archimedean 12
4.1. Norms on G 12
4.2. Inequalities regarding norms 13
4.3. Some spaces of rapidly decreasing functions on G 14
4.4. Realizing a generalized Whittaker model on $N\backslash G$ 17
5. Dual pairs and theta lifting of nilpotent orbits 19
5.1. Reductive dual pairs of type I 19
5.2. Moment maps and lifting of nilpotent orbits 20
5.3. Lifting of $\mathfrak{sl}_2$-triples 22
5.4. Isomorphism of $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ with a symplectic subspace $W_{\gamma,\tilde{\gamma}}$ of $\operatorname{Hom}(V, \tilde{V})$ 23
6. Generalized Whittaker models and Howe correspondence 24
6.1. The smooth oscillator-Heisenberg representation associated to $W_{\gamma,\tilde{\gamma}}$ 24
6.2. The main result and the key proposition 26
6.3. Strategy for the key proposition 27
6.4. The inductive step: relating covariants of $\mathscr{Y}_{(r),(r+1)}$ with $\mathscr{Y}_{(r-1),(r)}$ 31
6.5. From $\mathscr{Y}_{(r),(r+1)}$ to $\mathscr{H}_{\gamma,\tilde{\gamma}}$: proof of Proposition 6.4 38
Appendix A. Some facts on vector valued distributions 40

2000 *Mathematics Subject Classification.* 22E46 (Primary).
Key words and phrases. Local theta lifting, generalized Whittaker models, nilpotent orbits.
Both authors are supported by MOE2010-T2-2-113.

A.1. Vector valued distributions 40
A.2. Transverse order of a $\mathscr{V}$-distribution 41
A.3. Transverse jet bundle 41
A.4. Transverse symbols 42
References 42

1. Introduction and main result

Let $\Bbbk$ be a local field of characteristic 0, for which we fix a non-trivial unitary character ψ. Let G be a reductive group over $\Bbbk$, $\mathfrak{g}$ its Lie algebra, on which we fix an $\operatorname{Ad} G$-invariant non-degenerate bilinear form κ. Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ be an $\mathfrak{sl}_2$-triple, that is

$$[H, X] = 2X, \qquad [H, Y] = -2Y, \qquad [X, Y] = H.$$

Set $\mathfrak{g}_j = \{Z \in \mathfrak{g} \mid \operatorname{ad}(H)Z = jZ\}$, for $j \in \mathbb{Z}$. Then, from standard $\mathfrak{sl}_2$-theory, we have a finite direct sum $\mathfrak{g} = \oplus_{j\in\mathbb{Z}}\mathfrak{g}_j$. Define the Lie subalgebras $\mathfrak{u} = \oplus_{j\le -2}\mathfrak{g}_j$, $\mathfrak{n} = \oplus_{j\le -1}\mathfrak{g}_j$, $\mathfrak{p} = \oplus_{j\le 0}\mathfrak{g}_j$ and $\mathfrak{m} = \mathfrak{g}_0$. Let U, N, P, and M be the corresponding subgroups of G. Thus $U = \exp \mathfrak{u}$, $N = \exp \mathfrak{n}$, $P = \{p \in G \mid \operatorname{Ad}(p)\mathfrak{p} \subset \mathfrak{p}\}$ and $M = \{m \in G \mid \operatorname{Ad}(m)H = H\}$. Let

$$\chi_\gamma(\exp Z) = \psi(\kappa(X, Z)), \qquad \forall\ Z \in \mathfrak{u}. \tag{1.1}$$

Since $\kappa(X, [\mathfrak{u}, \mathfrak{u}]) = 0$, χ_γ defines a character on U. Let $M_X = \{m \in M \mid \operatorname{Ad}(m)X = X\}$. Then it is known [2, Section 3.4] that

$$M_X = G_\gamma := \{g \in G \mid \operatorname{Ad}(g)X = X,\ \operatorname{Ad}(g)Y = Y,\ \operatorname{Ad}(g)H = H\}\ .$$

In particular M_X is reductive. For the moment assume that $\mathfrak{g}_{-1} \neq 0$, or equivalently $\mathfrak{u} \subsetneq \mathfrak{n}$. In this case $\operatorname{ad}(X)|_{\mathfrak{g}_{-1}} : \mathfrak{g}_{-1} \longrightarrow \mathfrak{g}_1$ is an isomorphism, and we may define a symplectic structure on $\mathfrak{g}_{-1}$ by setting

$$\kappa_{-1}(S, T) = \kappa(\operatorname{ad}(X)S, T) = \kappa(X, [S, T]), \qquad \text{for all } S,\ T \in \mathfrak{g}_{-1}. \tag{1.2}$$

The associated Heisenberg group is $N/\operatorname{Ker}\chi_\gamma$, with center $U/\operatorname{Ker}\chi_\gamma$, where $\operatorname{Ker}\chi_\gamma$ denotes the kernel of χ_γ. Then, according to the Stone-von Neumann theorem, there exists a unique, up to equivalence, smooth irreducible (unitarizable) representation $(\rho_{\chi_\gamma}, \mathscr{S}_{\chi_\gamma})$ of N such that U acts by the character χ_γ. Since M_X preserves γ, it is well-known [31] that there exists a central cover of M_X, to be denoted by M_{χ_γ}, and a representation of a semi-direct product $M_{\chi_\gamma} \ltimes N$ on $\mathscr{S}_{\chi_\gamma}$ which extends the representation ρ_{χ_γ} of N. (Depending on the circumstances, this central cover may be taken to be either the group M_X itself, or a double cover.) We refer to the representation of $M_{\chi_\gamma} \ltimes N$ on $\mathscr{S}_{\chi_\gamma}$ as the smooth oscillator-Heisenberg representation. If $\mathfrak{g}_{-1} = 0$, then we define M_{χ_γ} to be just M_X. For notational convenience, we also denote by $(\rho_{\chi_\gamma}, \mathscr{S}_{\chi_\gamma})$ the 1-dimensional representation of $N = U$ given by the character χ_γ.

In this article, a smooth representation of a reductive group over $\Bbbk$ will mean a smooth representation in the usual sense for $\Bbbk$ non-Archimedean, namely it is locally constant, and a Casselman-Wallach representation for $\Bbbk$ Archimedean. The reader may consult [42, Chapter 11] for more information about Casselman-Wallach representations.

Definition. Fix an $\mathfrak{sl}_2$-triple $\gamma = \{X, H, Y\} \subset \mathfrak{g}$. Let $(\pi, \mathscr{V})$ be a smooth representation of G. We define the *space of generalized Whittaker models of* π *associated to* γ to be

$$\operatorname{Wh}_\gamma(\pi) = \operatorname{Hom}_N(\mathscr{V}, \mathscr{S}_{\chi_\gamma}). \tag{1.3}$$

Note that $\operatorname{Wh}_\gamma(\pi)$ is naturally an M_{χ_γ}-module.

From the well-known results of Jacobson-Morozov and Kostant [2, Chapter 3], the map $\gamma = \{X, H, Y\} \mapsto \mathcal{O} = \operatorname{Ad} G \cdot X$ yields a 1-1 correspondence between

$$\left\{ \begin{array}{c} \text{Conjugacy classes of} \\ \mathfrak{sl}_2\text{-triples in } \mathfrak{g} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Nilpotent } \operatorname{Ad} G\text{-orbits} \\ \mathcal{O} \subset \mathfrak{g} \end{array} \right\}.$$

If the conjugacy class of an $\mathfrak{sl}_2$-triple γ corresponds to a nilpotent orbit $\mathcal{O} \subset \mathfrak{g}$ we say that γ is an *$\mathfrak{sl}_2$-triple of type $\mathcal{O}$*. It is clear that given two conjugate $\mathfrak{sl}_2$-triples γ, γ', there exists an isomorphism $\phi : \mathrm{Wh}_\gamma(\pi) \longrightarrow \mathrm{Wh}_{\gamma'}(\pi)$ that intertwines the action of M_{χ_γ} and $M_{\chi_{\gamma'}}$. For this reason, sometimes we will abuse notation and denote $\mathrm{Wh}_\gamma(\pi)$ just by $\mathrm{Wh}_{\mathcal{O}}(\pi)$ and we will call it the *space of generalized Whittaker models associated to $\mathcal{O}$*, or the *space of generalized Whittaker models of type $\mathcal{O}$*. Note that this definition is (essentially) equivalent to the one given in [43, 23].

The study of Whittaker and generalized Whittaker models for representations of reductive groups over local fields evolved in connection with the theory of automorphic forms (via their Fourier coefficients), and has found important applications in both areas. See for example [34, 26, 15, 13, 43, 40], and a recent article of Jiang [12] discussing its role in a general theory of periods of automorphic forms. From the point of view of representation theory, the space of generalized Whittaker models may be viewed as one kind of nilpotent invariant associated to smooth representations. Another nilpotent invariant is the wave front cycle:

$$\mathrm{WF}(\pi) = \sum_{\substack{\mathcal{O} \subset \mathfrak{g} \\ \text{nilpotent}}} c_{\mathcal{O}}(\pi)[\mathcal{O}],$$

defined by Harish-Chandra in the non-Archimedean case [6] and by Barbasch and Vogan in the Archimedean case [1]. For $\mathfrak{k}$ non-Archimedean, Mœglin and Waldspurger [23] have established that $\mathrm{WF}(\pi)$ completely controls the spaces of generalized Whittaker models of interest, namely, if $\mathcal{O}$ is a nilpotent orbit which is maximal in the wave front set of π [10], then

$$c_{\mathcal{O}}(\pi) = \dim \mathrm{Wh}_{\mathcal{O}}(\pi).$$

For $\mathfrak{k}$ Archimedean, the corresponding phenomenon is not yet (fully) understood, except for the representations with the largest Gelfand-Kirillov dimension [37, 20] and unitary highest weight modules [44].

On the other hand, when $\mathfrak{k}$ is Archimedean, there is another nilpotent invariant defined by Vogan [38], which is called the associated cycle:

$$\mathrm{AC}(\pi) = \sum_{\substack{\mathcal{O} \subset \mathfrak{s} \\ \text{nilpotent}}} d_{\mathcal{O}}(\pi)[\mathcal{O}].$$

Here $\mathfrak{g}_{\mathbb{C}} = \mathfrak{k} \oplus \mathfrak{s}$ is the complexified Cartan decomposition. By the work of Schmid and Vilonen [32], we know that the wave front cycle and the associate cycle determine each other through the Kostant-Sekiguchi correspondence [33].

Now we come to the other part of the title regarding local theta lifting. Let $(G, \tilde{G})$ be a reductive dual pair in the sense of Howe [7]. Let $(\omega, \mathscr{Y})$ be the smooth oscillator representation associated to the dual pair $(G, \tilde{G})$ and to the character ψ of $\mathfrak{k}$. It is well-known that $(\omega, \mathscr{Y})$ yields a representation of $G \times \tilde{G}$ except when G is the isometry group of an odd dimensional quadratic space, in which case it is a representation of a double cover of $\tilde{G}$. In this exceptional case, we shall simply redefine $\tilde{G}$ to be the induced double cover, so as to simplify notation. Also in this case we shall only consider genuine representations of $\tilde{G}$, namely those representations such that the non-trivial element of the covering map acts by -1. For all other cases, representations are called genuine by convention.

Let $(\pi, \mathscr{V})$ be a smooth irreducible genuine representation of G, and consider the maximal π-isotypic quotient of $\mathscr{Y}$, called the Howe quotient of π. If it is non-zero, it is of the form $\mathscr{V} \otimes \Theta(\mathscr{V})$ (algebraic tensor product for $\mathfrak{k}$ non-Archimedean, and completed projective tensor product for $\mathfrak{k}$ Archimedean), where $\Theta(\mathscr{V})$ carries a smooth representation $\Theta(\pi)$ of $\tilde{G}$. The representation $\Theta(\pi)$ is sometimes referred to as the full theta lift (or colloquially the big theta lift) of π. Results of Howe [11] and Waldspurger [39] say that $\Theta(\pi)$ is an admissible representation of finite length, moreover, with the possible exception when $\mathfrak{k}$ is a dyadic field, $\Theta(\pi)$ has a unique irreducible quotient $\theta(\pi)$ called the (local) theta lift of π. In this article we shall focus on the full theta lift $\Theta(\pi)$, and shall thus place no restriction on the residue characteristic of $\mathfrak{k}$.

Let V, $\tilde{V}$, be the standard modules of G, $\tilde{G}$, respectively; and consider the moment maps φ, $\tilde{\varphi}$, from $\operatorname{Hom}(V, \tilde{V})$ to $\mathfrak{g}$ and $\tilde{\mathfrak{g}}$, respectively:

$$\begin{tikzcd}
 & \operatorname{Hom}(V, \tilde{V}) \arrow[dl, "\varphi"'] \arrow[dr, "\tilde{\varphi}"] & \\
\mathfrak{g} & & \tilde{\mathfrak{g}}
\end{tikzcd}$$

If $T \in \operatorname{Hom}(V, \tilde{V})$, then $\varphi(T)$ is nilpotent if and only if $\tilde{\varphi}(T)$ is nilpotent. As usual, this yields a notion of correspondence for nilpotent orbits $\mathcal{O} \subset \mathfrak{g}$ and $\tilde{\mathcal{O}} \subset \tilde{\mathfrak{g}}$. (In general this may not be one to one.) Denote by $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ the set of full rank elements in $\operatorname{Hom}(V, \tilde{V})$. Without any loss of generality, we assume that $\dim V \leq \dim \tilde{V}$, and elements of $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ are then represented by injective maps from V to $\tilde{V}$. In this article, we shall only be concerned with nilpotent orbits $\mathcal{O} \subset \mathfrak{g}$ in the image of $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ under the moment map φ, namely it satisfies

$$\varphi^{-1}(\mathcal{O}) \cap \operatorname{Max}\operatorname{Hom}(V, \tilde{V}) \neq \emptyset. \tag{1.4}$$

This will be our standing assumption. We write the resulting nilpotent orbit correspondence as $\mathcal{O} \mapsto \tilde{\mathcal{O}} = \Theta(\mathcal{O})$. See Section 5.2 for details.

As noted in the beginning, the space of generalized Whittaker models depends on an $\mathfrak{sl}_2$-triple. To relate the two $\mathfrak{sl}_2$-triples in $\mathfrak{g}$ and $\tilde{\mathfrak{g}}$, we make the following

Definition. Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\tilde{\mathcal{O}}$, respectively. We say that $T \in \operatorname{Hom}(V, \tilde{V})$ *lifts* γ to $\tilde{\gamma}$ if $\varphi(T) = X$, $\tilde{\varphi}(T) = \tilde{X}$ and $T(V_j) \subset \tilde{V}_{j+1}$ for all j. Here $V_j = \{v \in V \mid Hv = jv\}$, and likewise for $\tilde{V}_{j+1}$.

Set

$$\mathcal{O}_{\gamma,\tilde{\gamma}} = \{T \in \operatorname{Hom}(V, \tilde{V}) \mid T \text{ lifts } \gamma \text{ to } \tilde{\gamma}\},$$

and

$$\mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}} = \mathcal{O}_{\gamma,\tilde{\gamma}} \cap \operatorname{Max}\operatorname{Hom}(V, \tilde{V}). \tag{1.5}$$

(By Lemma 5.7, this is a single $M_X \times \tilde{M}_{\tilde{X}}$-orbit.)

Our main result, in a slightly less concrete form than what will be proved in Section 6, is

Theorem 1.1. *Let $(G, \tilde{G})$ be a reductive dual pair, and let $(\omega, \mathscr{Y})$ be the smooth oscillator representation associated to the dual pair $(G, \tilde{G})$ and to the character ψ of $\mathfrak{k}$. Let $(\pi, \mathscr{V})$ be a smooth irreducible genuine representation of G. Let $\mathcal{O} \subset \mathfrak{g}$ be a nilpotent G-orbit in the image of* $\operatorname{Max}\operatorname{Hom}(V, \tilde{V})$ *under the moment map φ and let $\tilde{\mathcal{O}} = \Theta(\mathcal{O}) \subset \tilde{\mathfrak{g}}$ be the corresponding nilpotent $\tilde{G}$-orbit. Then*

$$\mathrm{Wh}_{\Theta(\mathcal{O})}(\Theta(\pi)) \cong \mathrm{Wh}_{\mathcal{O}}(\check{\pi}),$$

where $\check{\pi}$ is the contragredient representation of π. More precisely, let γ and $\tilde{\gamma}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. Then, given any $T = T_{\gamma,\tilde{\gamma}} \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$, there is a surjective homomorphism

$\phi_T : \widetilde{M}_{\chi_{\tilde{\gamma}}} \twoheadrightarrow M_{\chi_\gamma}$ *(depending on* T*), inducing an action of* $\widetilde{M}_{\chi_{\tilde{\gamma}}}$ *on* $\mathrm{Wh}_\gamma(\check{\pi})$*, such that*

$$\mathrm{Wh}_{\tilde{\gamma}}(\Theta(\pi)) \cong \mathrm{Wh}_\gamma(\check{\pi})$$

as $\widetilde{M}_{\chi_{\tilde{\gamma}}}$*-modules.*

Remarks: (a) Nilpotent orbits in $\mathfrak{g}$ may be parameterized by equivalence classes of admissible ϵ-Hermitian Young tableaux (Section 3.2). Using this parametrization, the correspondence $\mathcal{O} \mapsto \Theta(\mathcal{O})$ can be described explicitly and in simple terms. Note that the assumption in (1.4) allows us to "embed" the sesquilinear Young tableau parameterizing $\mathcal{O}$ into the sesquilinear Young tableau parameterizing $\Theta(\mathcal{O})$. See Section 5.2.
(b) One particularly important circumstance is when $(G, \tilde{G})$ is in the stable range with G the smaller member ([8]; cf. [18]). Then every nilpotent orbit $\mathcal{O} \subset \mathfrak{g}$ is in the image of $\mathrm{Max}\,\mathrm{Hom}(V, \tilde{V})$ under the moment map φ, and for any smooth irreducible genuine representation $(\pi, \mathscr{V})$ of G, we have $\Theta(\pi) \neq 0$ [22, 28]. In this case and for $\mathfrak{k}$ non-Archimedean, Theorem 1.1 is due to Mœglin [21]. Our approach, explained in the following paragraph, is in some sense more conceptual. For the type of questions considered in this article, it is also well-known that substantially more effort is required to treat the Archimedean case.
(c) Assume that we are in the stable range and $\mathfrak{k}$ is Archimedean. There is a similar notion of theta lifting of nilpotent $K_{\mathbb{C}}$-orbits and one similarly expects a correspondence of associate cycles. We refer the reader to [44, 25] for some results in this direction, and the definitive result in the recent paper of Loke and Ma [19]. We also refer to earlier works of Przebinda [29, 30] on correspondence of wave front sets.
(d) Generically $\Theta(\pi)$ should be irreducible. One then obtains the space of generalized Whittaker models for $\theta(\pi)$. For example it is expected that under the assumption of stable range, $\Theta(\pi)$ is irreducible whenever π is unitarizable. For $\mathfrak{k}$ Archimedean, this is established in [19].
(e) As a direct consequence of Theorem 1.1, we have $\Theta(\pi) \neq 0$, if $\mathrm{Wh}_{\mathcal{O}}(\check{\pi}) \neq 0$, for some $\mathcal{O}$ satisfying the condition in (1.4).

The main ingredient in the proof of Theorem 1.1 is our description of the space of covariants $\mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}}$ of the smooth oscillator representation $\mathscr{Y}$, where $(\tilde{U}, \chi_{\tilde{\gamma}})$ for the $\mathfrak{sl}_2$-triple $\tilde{\gamma}$ in $\tilde{\mathfrak{g}}$ is as (U, χ_γ) for the $\mathfrak{sl}_2$-triple γ, and $\tilde{\gamma}$ is a lift of γ. (This is reminiscent of the computation of the Jacquet modules of the smooth oscillator representation by Kudla [17], though not in method.) We show it is isomorphic to a certain space $\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ of rapidly decreasing functions on $N\backslash G$ with values in $\mathscr{H}_{\gamma,\tilde{\gamma}}$, where $\mathscr{H}_{\gamma,\tilde{\gamma}}$ is the space of the smooth oscillator-Heisenberg representation associated to a certain symplectic subspace $W_{\gamma,\tilde{\gamma}}$ of $\mathrm{Hom}(V, \tilde{V})$. This is the key Proposition 6.4. An important point is that we may construct an isomorphism of $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ with $W_{\gamma,\tilde{\gamma}}$, which depends on $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$. The key proposition basically says that we can define a natural surjective (matrix coefficient) map (using "homogeneous components" of T), from $\mathscr{Y}$ to $\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$, which induces an isomorphism on $\mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}}$!

Here are some additional words on the organization and contents of this article. In Section 2, we describe classical groups as the isometry groups of ϵ-Hermitian D-modules, where D is a division algebra over $\mathfrak{k}$. In Section 3, we review the well-known parametrization of nilpotent orbits in the classical Lie algebras, following the book by Collingwood and McGovern [2]. We also introduce the generalized Whittaker models associated to the nilpotent orbits. In Section 4, we introduce some Fréchet spaces of functions on G to realize the space of generalized Whittaker models for $\mathfrak{k}$ Archimedean. In Section 5, we recall the notion of lifting of nilpotent orbits in the setting of dual pairs (via the moment maps), and we describe the fine structure of lifting for those orbits $\mathcal{O}$ and $\tilde{\mathcal{O}}$ which correspond via injective maps from V to $\tilde{V}$. We also prove an explicit isomorphism of

$\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ with a symplectic subspace $W_{\gamma,\tilde{\gamma}}$ of $\mathrm{Hom}(V, \tilde{V})$. In Section 6, we relate the generalized Whittaker models of $\breve{\pi}$ and $\Theta(\pi)$. As mentioned previously, its main ingredient is the description of the space of covariants $\mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}}$ of the smooth oscillator representation $\mathscr{Y}$. To arrive at this, we make extensive use of the gradation in the standard modules V and $\tilde{V}$ given by the neutral elements H, $\tilde{H}$ of the two $\mathfrak{sl}_2$-triples $\gamma \subset \mathfrak{g}$ and $\tilde{\gamma} \subset \tilde{\mathfrak{g}}$ of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. On the one hand, it gives rise to totally isotropic subspaces and thus convenient realizations of $\mathscr{Y}$. On the other hand, it facilitates an inductive argument based on the heights of the gradations. Together with the isomorphism of $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ with $W_{\gamma,\tilde{\gamma}}$, this implies the relationship between the generalized Whittaker models of $\breve{\pi}$ and $\Theta(\pi)$.

Acknowledgements: the authors thank W. T. Gan, D. Jiang, and B. Sun for their interests and comments.

2. Classical groups as isometry groups of Hermitian modules

2.1. **Hermitian D-modules.** Let $\Bbbk$ be a local field, $|\cdot|$ its absolute value, and let ψ be a fix non-trivial unitary character of $\Bbbk$. Let D be one of the following division algebras over $\Bbbk$: the field $\Bbbk$ itself, a quadratic field extension of $\Bbbk$ or the quaternion division $\Bbbk$-algebra. Observe that D comes equipped with a canonical involutive anti-automorphism (the identity map, the non-trivial Galois element, or the main involution, respectively) which we will denote by $x \mapsto \overline{x}$. Throughout this article, we will only consider finitely generated modules over D.

Let V and W be two right D-modules. We will denote the set of right D-module morphisms from V to W by

$$\mathrm{Hom}_D(V, W) = \{T : V \longrightarrow W \,|\, T(v_1 a + v_2 b) = T(v_1)a + T(v_2)b \text{ for all } v_1,\, v_2 \in V,\, a,\, b \in D\}.$$

If $V = W$, we will denote this set by $\mathrm{End}_D(V)$. Set

$$\mathrm{GL}(V, D) = \{T \in \mathrm{End}_D(V) \,|\, T \text{ is invertible}\}.$$

When it is clear from the context what the division algebra D is, we may just omit D in various of these notations.

Let V' be the set of right D-linear functionals on V. There is a natural left D-module structure on V' given by setting

$$(a\lambda)(v) = a\lambda(v), \quad \text{for all } a \in D,\, v \in V, \text{ and } \lambda \in V'.$$

Observe that with this structure, $W \otimes_D V'$ is naturally isomorphic to $\mathrm{Hom}_D(V, W)$ as a $\Bbbk$-vector space. Given $T \in \mathrm{Hom}_D(V, W)$, we will specify an element in $\mathrm{Hom}_D(W', V')$ (analogously defined), which we will also denote T, by setting $(\lambda T)(v) := \lambda(Tv)$, for $v \in V$ and $\lambda \in W'$. This correspondence gives rise to natural isomorphisms between $\mathrm{End}_D(V)$ and $\mathrm{End}_D(V')$, and between $\mathrm{GL}_D(V)$ and $\mathrm{GL}_D(V')$.

Definition 2.1. Let $\varepsilon = \pm 1$. We say that (V, B) is a right ε-Hermitian D-module if V is a right D-module and B is an ε-Hermitian form, i.e., $B : V \times V \longrightarrow D$ is a map such that

(1) B is *sesquilinear*: for all v_1, v_2, $v_3 \in V$, a, $b \in D$,

$$B(v_1, v_2 a + v_3 b) = B(v_1, v_2)a + B(v_1, v_3)b \quad \text{and} \quad B(v_1 a + v_2 b, v_3) = \overline{a}B(v_1, v_3) + \overline{b}B(v_2, v_3).$$

(2) B is *ε-Hermitian*:

$$B(v, w) = \varepsilon\overline{B(w, v)} \qquad \text{for all } v, w \in V.$$

(3) B is *non-degenerate*.

Given a right ε-Hermitian D-module (V,B), we may construct a left ε-Hermitian D-module (V^*,B^*) in the following way: as a set, V^* will be the set of symbols $\{v^* \mid v \in V\}$. Then we give to V^* a left D-module structure by setting, for all v, $w \in V$, $a \in D$,

$$v^* + w^* = (v+w)^* \text{ and } av^* = (v\overline{a})^*.$$

Finally, we set

$$B^*(v^*,w^*) = \overline{B(w,v)} \qquad \text{for all } v,\, w \in V.$$

In an analogous way, we may define the $*$ operation on left ε-Hermitian D-modules. Then V^{**} is naturally isomorphic with V. Given $T \in \operatorname{End}_D(V)$, we define $T^* \in \operatorname{End}_D(V^*)$ by setting $v^*T^* := (Tv)^*$. With this definition, it is easily seen that $(TS)^* = S^*T^*$, for all S, $T \in \operatorname{End}_D(V)$. Therefore the map $g \mapsto (g^*)^{-1}$ defines a group isomorphism between $\mathrm{GL}_D(V)$ and $\mathrm{GL}_D(V^*)$.

Observe that the form B induces a left D-module isomorphism $B^\flat : V^* \longrightarrow V'$ given by $B^\flat(v^*)(w) = B(v,w)$ for v, $w \in V$. In what follows, we will make implicit use of this map to identify these two spaces. With this identification, for any $T \in \operatorname{End}_D(V)$, we can think of T^* as an element in $\operatorname{End}_D(V)$ defined by $v^*(T^*w) := (v^*T^*)(w)$, i.e., T^* is defined by the usual condition that

$$B(v,T^*w) = B(Tv,w) \qquad \text{for all } v,\, w \in V.$$

A D-submodule $E \subset V$ is said to be *totally isotropic* if $B|_{E\times E} = 0$. If E is a totally isotropic submodule, then there exists a totally isotropic submodule $F \subset V$ such that $B|_{(E\oplus F)\times(E\oplus F)}$ is non-degenerate. If we set

$$U = (E \oplus F)^\perp := \{u \in V \mid B(u,w) = 0 \text{ for all } w \in E \oplus F\},$$

then $V = E \oplus F \oplus U$, and $B|_{U\times U}$ is non-degenerate. In this case we say that E and F are totally isotropic, *complementary* submodules. Observe that then $B^\flat|_{F^*} : F^* \longrightarrow E'$ is an isomorphism. As before we will make implicit use of this isomorphism to identify F^* with E'.

2.2. **Isometry groups.** Given a right ε-Hermitian D-module (V,B), we define its isometry group

$$G(V,B) = \{g \in \mathrm{GL}(V) \mid B(gv,gw) = B(v,w) \text{ for all } v,\, w \in V\}.$$

When there is no risk of confusion regarding B, we will denote this group just by $G(V)$ or even just as G. Observe that if $g \in G$, then $g^* = g^{-1}$.

Associated to the group $\mathrm{GL}(V)$ we have the Lie algebra $\mathfrak{gl}(V) = \operatorname{End}(V)$ with bracket $[T,S] := TS - ST$, for all T, $S \in \mathfrak{gl}(V)$. Similarly, associated to the group G we have the Lie algebra

$$\begin{aligned} \mathfrak{g} &= \{T \in \mathfrak{gl}(V) \mid B(Tv,w) + B(v,Tw) = 0 \text{ for all } v,\, w \in V\} \\ &= \{T \in \mathfrak{gl}(V) \mid T^* = -T\}. \end{aligned}$$

We define a bilinear form on $\mathfrak{g}$ by

$$\kappa(T,S) = \operatorname{Tr}(T^*S)/2,$$

for all T, $S \in \mathfrak{g}$. Here $\operatorname{Tr}(T^*S)$ is the trace of T^*S as a $\mathfrak{k}$-linear transformation. This bilinear form is non-degenerate and *invariant* with respect to the adjoint action of G on $\mathfrak{g}$.

Remark 2.2. In the literature, the isometry groups $G(V,B)$ are called type I. Given a division algebra D, one may consider the algebra $\mathbb{D} = D \oplus D$, which has the canonical involution $\overline{(a,b)} = (b,a)$, for a, $b \in D$. Then one may similarly define right ϵ-Hermitian $\mathbb{D}$-modules ($\epsilon = \pm 1$) and their isometry groups which are easily seen to be isomorphic to $\mathrm{GL}(V,D)$, where V are right D-modules. They are called type II. With this modification (from D to $\mathbb{D}$), all results in this article, which are stated for type I dual pairs, carry over to the case of type II dual pairs.

3. NILPOTENT ORBITS OF CLASSICAL GROUPS

3.1. **Nilpotent orbits and $\mathfrak{sl}_2$-triples.** In order to set up notation, and facilitate the exposition of the reminder of this article, we will review the basic structural results on nilpotent orbits and $\mathfrak{sl}_2$-triples in $\mathfrak{g}$. The exposition given here is based on the book of Collingwood and McGovern [2].

Let $X \in \mathfrak{g}$ be a nilpotent element. Then, by the Jacobson-Morozov theorem, there exists an $\mathfrak{sl}_2$-triple $\gamma = \{H, X, Y\} \subset \mathfrak{g}$, containing X as the nilpositive element, namely

$$[H, X] = 2X, \qquad [H, Y] = -2Y, \qquad [X, Y] = H.$$

Let $\mathfrak{g}_i = \{Z \in \mathfrak{g} \mid \mathrm{ad}(H)(Z) = iZ\}$, for $i \in \mathbb{Z}$. Then, from standard $\mathfrak{sl}_2$-theory, we have that

$$\mathfrak{g} = \bigoplus_{i\in\mathbb{Z}} \mathfrak{g}_i. \tag{3.1}$$

Now let

$$\mathfrak{p} = \bigoplus_{i\le 0} \mathfrak{g}_i, \qquad \overline{\mathfrak{p}} = \bigoplus_{i\ge 0} \mathfrak{g}_i,$$

and set

$$P = \{g \in G \mid \mathrm{Ad}(g)\mathfrak{p} \subset \mathfrak{p}\}, \qquad \overline{P} = \{g \in G \mid \mathrm{Ad}(g)\overline{\mathfrak{p}} \subset \overline{\mathfrak{p}}\}.$$

$\overline{P}$ (resp. $\overline{\mathfrak{p}}$) is called the Jacobson-Morozov parabolic subgroup (resp. parabolic subalgebra) associated to X. Observe that P (resp. $\mathfrak{p}$) is a parabolic subgroup opposite to $\overline{P}$ (resp. parabolic subalgebra opposite to $\overline{\mathfrak{p}}$). (The subgroup $\overline{P}$ depends only on X, but the subgroup P depends on the choice of $\mathfrak{sl}_2$-triple γ containing X.) Let $M = \{m \in G \mid \mathrm{Ad}(m)H = H\}$. Its Lie algebra is $\mathfrak{m} = \{Z \in \mathfrak{g} \mid \mathrm{ad}(Z)H = 0\}$, which is exactly $\mathfrak{g}_0$. Let

$$\mathfrak{n} = \bigoplus_{i\le -1} \mathfrak{g}_i \qquad \text{and} \qquad \mathfrak{u} = \bigoplus_{i\le -2} \mathfrak{g}_i.$$

Observe that if $Z \in \mathfrak{n}$, then Z is nilpotent and hence $\exp Z = \sum_{j=0}^{\infty} Z^j/j!$ is a well defined element in $\mathrm{End}(V)$. Let $N = \exp \mathfrak{n} = \{\exp Z \mid Z \in \mathfrak{n}\}$ and $U = \exp \mathfrak{u} = \{\exp Z \mid Z \in \mathfrak{u}\}$; then U, N are subgroups of G, and $P = MN$. Similarly we have the Levi decomposition $\overline{P} = M\overline{N}$, with $\overline{N}$ the unipotent radical of $\overline{P}$.

For $i \in \mathbb{Z}$, let $V_i = \{v \in V \mid Hv = iv\}$. Then, again from standard $\mathfrak{sl}_2$-theory,

$$V = \bigoplus_{i\in\mathbb{Z}} V_i. \tag{3.2}$$

We may also characterize M as the set of $m \in G$ that preserves the direct sum decomposition given in (3.2). Now observe that, since $H^* = -H$, $B|_{V_0\times V_0}$ is non-degenerate and B establishes a perfect pairing between V_i and V_{-i} for all $i > 0$. Using this pairing we define for any $T \in \mathrm{End}(V_i)$ a map $T^* \in \mathrm{End}(V_{-i})$ given by

$$B(Tv, w) = B(v, T^*w), \qquad \text{for all } v \in V_i,\ w \in V_{-i}.$$

It follows immediately that, if $g \in \mathrm{GL}(V_i)$, then $B(gv, (g^*)^{-1}w) = B(v, w)$, for all $v \in V_i$, $w \in V_{-i}$. From this we conclude that there is an embedding of $\mathrm{GL}(V_i)$ into M for all $i > 0$. We will denote the image of this embedding by M_i. Proceeding in a similar manner we may also define a natural embedding of $G(V_0, B|_{V_0\times V_0})$ into M, whose image we will denote by M_0. Then

$$M = \prod_{i\ge 0} M_i \cong G(V_0) \times \prod_{i>0} \mathrm{GL}(V_i). \tag{3.3}$$

Set $\mathfrak{g}_\gamma = \mathrm{Span}_{\Bbbk}\{X, H, Y\} \subset \mathfrak{g}$. We have the $\mathfrak{g}_\gamma$-isotypic decomposition

$$V = \bigoplus_{j=1}^{l} V^{\gamma,t_j}, \tag{3.4}$$

where V^{γ,t_j} is a direct sum of irreducible t_j-dimensional $\mathfrak{g}_\gamma$-modules, and $t_1 > t_2 > \ldots > t_l > 0$. Let $V_i^{\gamma,t_j} = V^{\gamma,t_j} \cap V_i$. It is nonzero if and only if $|i| < t_j$ and $i \equiv t_j - 1 \pmod 2$. Then it is clear that

$$V_i = \bigoplus_{j,\, t_j > |i|} V_i^{\gamma,t_j}. \tag{3.5}$$

Using standard results from the representation theory of $\mathfrak{sl}_2$, we have that for all $i \geq 0$, the map $(X|_{V_{-i}})^i : V_{-i} \longrightarrow V_i$ is invertible. (Here we are using the convention that $(X|_{V_0})^0$ is the identity map on V_0.) The statement remains true for $i < 0$ if we interpret a negative power of an invertible map as the positive power of its inverse.

Now define a Hermitian form B_i on V_i by setting $B_i(v, w) = B(X|_{V_i}^{-i} v, w)$ for all v, $w \in V_i$. Since B establishes a perfect pairing between V_i and V_{-i}, it is clear that B_i is non-degenerate. The analysis applied to (V, B) applies equally to $(V^{\gamma,t_j}, B^{\gamma,t_j})$, where B^{γ,t_j} is the restriction of B to V^{γ,t_j}. Thus B_i is in fact non-degenerate when restricted to any V_i^{γ,t_j}. We will denote by B_i^{γ,t_j} the restriction of B_i to V_i^{γ,t_j}. The following result is also clear:

(3.6)

$$\text{If } -t_j + 1 \leq i < i + 2 \leq t_j - 1, \qquad \text{then } B_{i+2}(Xv, Xw) = -B_i(v, w) \qquad \text{for all } v,\ w \in V_i^{\gamma,t_j}.$$

In particular, all the B_i^{γ,t_j}'s are determined by $B_{t_j-1}^{\gamma,t_j}$ (on $V_{t_j-1}^{\gamma,t_j}$, the space of highest weight vectors in V^{γ,t_j}).

Let $M_X = \{m \in M \mid \mathrm{Ad}(m)X = X\}$. The following result is due to Kostant.

Theorem. ([2, Section 3.4]) *Let* $G_X = \{g \in G \,|\, \mathrm{Ad}(g)X = X\}$. *Then*

$$G_X = M_X \overline{N}_X,$$

where $\overline{N}_X = \{n \in \overline{N} \mid \mathrm{Ad}(n)X = X\}$. *Furthermore*

$$M_X = G_\gamma := \{g \in G \,|\, \mathrm{Ad}(g)X = X,\ \mathrm{Ad}(g)H = H,\ \mathrm{Ad}(g)Y = Y\}.$$

We now have all the ingredients to give a description of M_X in the spirit of the description of M given in equation (3.3). Observe that M_X acts on $V_{t_j-1}^{\gamma,t_j}$ preserving $B_{t_j-1}^{\gamma,t_j}$. From this observation and equation (3.6) we conclude that there is an embedding of $G(V_{t_j-1}^{\gamma,t_j}, B_{t_j-1}^{\gamma,t_j})$ into M_X. Let M_{X,t_j-1} be the image of this embedding. Then

$$M_X = \prod_{j=1}^{l} M_{X,t_j-1} \cong \prod_{j=1}^{l} G(V_{t_j-1}^{\gamma,t_j}). \tag{3.7}$$

3.2. **ϵ-Hermitian Young tableaux.** A partition of n is a tuple $[d_1, \ldots, d_k]$ of positive integers such that $d_1 \geq d_2 \geq \ldots \geq d_k$ and $d_1 + \cdots + d_k = n$. Partitions of n are frequently represented by *Young diagrams* in the following way: given a partition $\mathbf{d} = [d_1, \ldots, d_k]$ we construct a left-justified array of empty boxes such that the i-th row has d_i boxes. This array is the Young diagram associated to $\mathbf{d}$. Another way of describing a partition $\mathbf{d} = [d_1, \ldots, d_k]$ of n is using the *exponential notation* $\mathbf{d} = [t_1^{i_1}, \ldots, t_l^{i_l}]$, which means that the number t_k appears i_k times (the multiplicity) in the partition, and $t_1 > \cdots > t_l > 0$.

Definition 3.1. A *sesquilinear Young tableau* is a pair $\Gamma = (\mathbf{d}^\Gamma, (V^\Gamma, B^\Gamma))$ such that $\mathbf{d}^\Gamma = [t_1^{i_1}, \ldots, t_l^{i_l}]$ is a partition of n, and (V^Γ, B^Γ) is an assignment, for each $1 \leq j \leq l$, of an ϵ_j-Hermitian module $(V_{t_j-1}^{\Gamma,t_j}, B_{t_j-1}^{\Gamma,t_j})$ of dimension i_j.

Example 3.2. The following picture represents a sesquilinear Young tableau $\Gamma = (\mathbf{d}^\Gamma, (V^\Gamma, B^\Gamma))$, where $\mathbf{d}^\Gamma = [3^2, 2^3, 1^2]$ is a partition of 14, and $V_2^{\Gamma,3}$, $V_1^{\Gamma,2}$, $V_0^{\Gamma,1}$ have dimensions 2, 3 and 2, respectively.


$(V_2^{\Gamma,3}, B_2^{\Gamma,3})$
$(V_1^{\Gamma,2}, B_1^{\Gamma,2})$
$(V_0^{\Gamma,1}, B_0^{\Gamma,1})$

Let $(\rho_m, \Bbbk^m)$ be the irreducible representation of $\mathfrak{sl}_2$ of dimension m, and fix, for each m, a non-degenerate invariant bilinear form $(\cdot,\cdot)_m$ on $\Bbbk^m$. Recall that this form is unique up to scalar and that $(f_1, f_2)_m = (-1)^{m-1}(f_2, f_1)_m$, for all f_1, $f_2 \in \Bbbk^m$. This follows from the analysis surrounding equation (3.6).

Given a sesquilinear Young tableau $\Gamma = (\mathbf{d}^\Gamma, (V^\Gamma, B^\Gamma))$ we will set

$$V^{\Gamma,t_j} := V_{t_j-1}^{\Gamma,t_j} \otimes_{\Bbbk} \Bbbk^{t_j}. \tag{3.8}$$

On this space we define a $(-1)^{t_j-1}\epsilon_j$-Hermitian form B^{Γ,t_j} by

$$B^{\Gamma,t_j}(v_1 \otimes f_1, v_2 \otimes f_2) = B_{t_j-1}^{\Gamma,t_j}(v_1, v_2)(f_1, f_2)_{t_j}, \qquad v_1,\ v_2 \in V_{t_j-1}^{\Gamma,t_j}, \text{ and } f_1,\ f_2 \in \Bbbk^{t_j}. \tag{3.9}$$

Definition 3.3. Let $\Gamma = (\mathbf{d}^\Gamma, (V^\Gamma, B^\Gamma))$ and $\Phi = (\mathbf{d}^\Phi, (V^\Phi, B^\Phi))$ be two sesquilinear Young Tableaux.

(i) We say that Γ and Φ are *equivalent* if $\mathbf{d}^\Gamma = \mathbf{d}^\Phi = [t_1^{i_1}, \ldots, t_l^{i_l}]$ and $(V_{t_j-1}^{\Gamma,t_j}, B_{t_j-1}^{\Gamma,t_j})$ is isomorphic to $(V_{t_j-1}^{\Phi,t_j}, B_{t_j-1}^{\Phi,t_j})$ for all $j = 1, \ldots, l$.

(ii) We say that Γ is *ϵ-Hermitian* if $(V^{\Gamma,t_j}, B^{\Gamma,t_j})$ is an ϵ-Hermitian module for all $j = 1, \ldots, l$.

(iii) Given an ϵ-Hermitian module (V, B), we say that Γ is *admissible* for (V, B), or (V, B)-*admissible*, if $(\oplus_j V^{\Gamma,t_j}, \oplus_j B^{\Gamma,t_j})$ is isomorphic to (V, B).

Recall that given an $\mathfrak{sl}_2$-triple $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ there exists $t_1 > \ldots > t_l$ such that $V = \oplus_j V^{\gamma,t_j}$. Using this decomposition we may define an ϵ-Hermitian Young tableaux $\Gamma_\gamma = (\mathbf{d}^{\Gamma_\gamma}, (V^{\Gamma_\gamma}, B^{\Gamma_\gamma}))$ by setting $\mathbf{d}^{\Gamma_\gamma} = [t_1^{i_1}, \ldots, t_l^{i_l}]$, where $i_j = \dim V_{t_j-1}^{\gamma,t_j}$, and

$$(V_{t_j-1}^{\Gamma_\gamma,t_j}, B_{t_j-1}^{\Gamma_\gamma,t_j}) := (V_{t_j-1}^{\gamma,t_j}, B_{t_j-1}^{\gamma,t_j}), \tag{3.10}$$

for all $j = 1, \ldots, l$.

Methods of [2, Section 9.3] imply that this assignment gives a bijection between the set of $\mathfrak{sl}_2$-triples in $\mathfrak{g}$ up to the Adjoint action of G and equivalence classes of admissible ϵ-Hermitian Young tableaux. We thus have

Proposition 3.4. *There is a* 1-1 *correspondence between the following sets:*

$$\{\textit{Nilpotent orbits in } \mathfrak{g}\} \longleftrightarrow \left\{ \begin{array}{c} \textit{Equivalence classes of admissible} \\ \epsilon\textit{-Hermitian Young tableaux} \end{array} \right\}.$$

Remark 3.5. If $\Bbbk$ is the field of real numbers then Hermitian Young tableaux can be more concretely described by *signed* Young diagrams [2, Section 9.3].

3.3. **Generalized Whittaker models associated to nilpotent orbits.** Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ be an $\mathfrak{sl}_2$-triple. As in the introduction, we define the character χ_γ of U, by $\chi_\gamma(\exp Z) = \psi(\kappa(X, Z))$ for all $Z \in \mathfrak{u}$; and a symplectic structure on $\mathfrak{g}_{-1}$ by setting

$$\kappa_{-1}(S,T) = \kappa(\mathrm{ad}(X)S, T) = \kappa(X, [S,T]) \qquad \text{for all } S,\ T \in \mathfrak{g}_{-1}.$$

(Note the similarity between this definition and the definition of the forms B_i on V_i in Section 3.1.) Let $\mathcal{H}_\gamma$ be the Heisenberg group associated to the symplectic space $(\mathfrak{g}_{-1}, \kappa_{-1})$. That is $\mathcal{H}_\gamma = \mathfrak{g}_{-1} \times \mathfrak{k}$, $\{0\} \times \mathfrak{k}$ is central, and $(T, 0)(S, 0) = (T + S, \kappa(T, S)/2)$ for all T, $S \in \mathfrak{g}_{-1}$. Then, according to the Stone-von Neumann theorem, there exists a unique, up to equivalence, *smooth* irreducible (unitarizable) representation $(\rho_\gamma, \mathscr{S}_\gamma)$ of H_γ such that the center of H_γ acts by the character ψ. Here smooth means that it is locally constant if $\mathfrak{k}$ is non-Archimedian, and if $\mathfrak{k}$ is Archimedian, $(\rho_\gamma, \mathscr{S}_\gamma)$ is the smoothing of the usual irreducible unitary representation of H_γ with the central character ψ.

Let $\alpha_\gamma : U \longrightarrow \mathfrak{k}$ be the map given by $\alpha_\gamma(\exp Z) = \kappa(X, Z)$, for all $Z \in \mathfrak{u}$. It is standard to check that α_γ defines a surjective group homomorphism. Furthermore, it extends to a group homomorphism $\alpha_\gamma : N \mapsto H_\gamma$ given by

$$\alpha_\gamma(\exp T \exp Z) = (T, \kappa(X, Z)), \qquad \text{for all } T \in \mathfrak{g}_{-1},\ Z \in \mathfrak{u}. \tag{3.11}$$

By composition, this yields a representation $(\rho_{\chi_\gamma}, \mathscr{S}_{\chi_\gamma})$ of N, where $\mathscr{S}_{\chi_\gamma} := \mathscr{S}_\gamma$ and

$$\rho_{\chi_\gamma}(n)v = \rho_\gamma(\alpha_\gamma(n))v, \qquad \text{for all } n \in N,\ v \in \mathscr{S}_{\chi_\gamma}. \tag{3.12}$$

Observe that then, for all $Z \in \mathfrak{u}$, $v \in \mathscr{S}_{\chi_\gamma}$,

$$\rho_{\chi_\gamma}(\exp Z)v = \rho_\gamma(0, \kappa(X, Z))v = \psi(\kappa(X, Z)) = \chi_\gamma(\exp Z).$$

In particular, if $\mathfrak{g}_{-1} = 0$, then $N = U$ acts on the 1-dimensional space $\mathscr{S}_{\chi_\gamma}$ by the character χ_γ. Since M_X preserves γ, it is well-known [31] that there exists a central cover of M_X, to be denoted by M_{χ_γ}, and a representation of a semi-direct product $M_{\chi_\gamma} \ltimes N$ on $\mathscr{S}_{\chi_\gamma}$ which extends the representation ρ_{χ_γ} of N. (Depending on the circumstances, this central cover may be taken to be either the group M_X itself, or a double cover.) We refer to the representation $(\rho_{\chi_\gamma}, \mathscr{S}_{\chi_\gamma})$ of $M_{\chi_\gamma} \ltimes N$ as the smooth oscillator-Heisenberg representation associated to χ_γ. We remark that there is a notion of "smooth Fréchet representations of moderate growth" for groups of the type $M_{\chi_\gamma} \ltimes N$. See [4, Definition 1.4.1] or [35, Section 2].

Definition 3.6. Let $(\pi, \mathscr{V})$ be a smooth representation of G, and let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ be an $\mathfrak{sl}_2$-triple. We define the *space of generalized Whittaker models of* π *associated to* γ to be

$$\mathrm{Wh}_\gamma(\pi) = \mathrm{Hom}_N(\mathscr{V}, \mathscr{S}_{\chi_\gamma}). \tag{3.13}$$

Note that $\mathrm{Wh}_\gamma(\pi)$ is naturally an M_{χ_γ}-module.

We also make the following definition.

Definition 3.7. Let $\mathcal{O} \subset \mathfrak{g}$ be a nilpotent orbit. We say that $\gamma = \{X, H, Y\}$ is an $\mathfrak{sl}_2$*-triple of type* $\mathcal{O}$ if $X \in \mathcal{O}$, where X is the nilpositive element of the $\mathfrak{sl}_2$-triple.

From the well-known results of Jacobson-Morozov and Kostant [2, Chapter 3], the map $\gamma = \{X, H, Y\} \mapsto \mathcal{O} = \mathrm{Ad}\, G \cdot X$ yields a 1-1 correspondence between

$$\left\{ \begin{array}{c} \text{Conjugacy classes of} \\ \mathfrak{sl}_2\text{-triples in } \mathfrak{g} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Nilpotent Ad}\, G\text{-orbits} \\ \mathcal{O} \subset \mathfrak{g} \end{array} \right\}.$$

As noted in the introduction, it is clear that given two conjugate $\mathfrak{sl}_2$-triples γ, γ', there will be an obvious isomorphism $\phi : \mathrm{Wh}_\gamma(\pi) \longrightarrow \mathrm{Wh}_{\gamma'}(\pi)$ that intertwines the action of M_{χ_γ} and $M_{\chi_{\gamma'}}$. By abuse of notation, we will denote $\mathrm{Wh}_\gamma(\pi)$ just by $\mathrm{Wh}_{\mathcal{O}}(\pi)$ and we will call it the *space of generalized Whittaker models associated to* $\mathcal{O}$, or the *space of generalized Whittaker models of type* $\mathcal{O}$.

4. A realization of generalized Whittaker models: $\mathfrak{k}$ Archimedean

Let $(\pi, \mathscr{V})$ be a smooth representation of G, and let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ be an $\mathfrak{sl}_2$-triple. In this section we will give convenient realizations of the spaces $\mathscr{S}_{\chi_\gamma}$ and $\mathrm{Wh}_\gamma(\pi)$ associated to γ and π. As we will see later in this section, the analysis required for $\mathfrak{k}$ Archimedean is substantially more involved than the case where $\mathfrak{k}$ is non-Archimedian. For this reason, we will devote this section to the Archimedean case and will be contented to just indicate how the analog results work in the non-Archimedian case.

4.1. **Norms on G.** Assume that $\mathfrak{k} = \mathbb{R}$ or $\mathbb{C}$. Let (V, B_V) be an ϵ-Hermitian module of dimension n. Then we have the following possibilities for $G = G(V)$:

(1) $\mathfrak{k} = \mathbb{R}$ and $D = \mathbb{R}$. In this case, either

$$G \cong O(p,q),\ p+q=n, \text{ if } \epsilon = 1, \qquad \text{or} \qquad G \cong Sp(n,\mathbb{R}) \text{ if } \epsilon = -1 \text{ and } n \text{ is even.}$$

(2) $\mathfrak{k} = \mathbb{R}$ and $D = \mathbb{C}$. In this case

$$G \cong U(p,q),\ p+q=n, \qquad\qquad \text{regardless of } \ \epsilon \ \ (= \pm 1).$$

(3) $\mathfrak{k} = \mathbb{R}$ and $D = \mathbb{H}$. In this case either

$$G \cong Sp(p,q),\ p+q=n, \text{ if } \epsilon = 1, \qquad \text{or} \qquad G \cong O^*(2n), \text{ if } \epsilon = -1.$$

(4) $\mathfrak{k} = \mathbb{C}$ and $D = \mathbb{C}$. Then

$$G \cong O(n,\mathbb{C}) \text{ if } \epsilon = 1 \qquad \text{or} \qquad G \cong Sp(n,\mathbb{C}) \text{ if } \epsilon = -1 \text{ and } n \text{ is even.}$$

Let $V = E \oplus U \oplus F$, with E and F totally isotropic, complementary submodules of maximal dimension. Thus U is anisotropic. Let $P_E = \mathrm{Stab}_E$, the stabilizer of E. Then $P_E = M_E N_E$, is a Langlands decomposition of P_E, where $M_E = \{m \in P_E \mid m \cdot F \subset F\}$ and the Lie algebra of N_E is

$$\mathfrak{n}_E \cong \mathrm{Hom}(U, E) \oplus \mathfrak{z},$$

with $\mathfrak{z} \cong \{Z : F \to E \mid Z^* = -Z\}$. It follows from this description that $M_E \cong \mathrm{GL}(E) \times G(U)$ and that $G(U)$ is compact. Thus there exists a real inner product B_U^+ on U such that $G(U,B) \subset G(U, B_U^+)$.

Let $\{e_1, \ldots, e_l\}$, and $\{f_1, \ldots, f_l\}$ be basis of E and F, respectively, such that

$$B_V(e_i, f_j) = \delta_{i,l-j+1}, \qquad \text{for all } 1 \leq i,j \leq l.$$

Then we can extend B_U^+ to an inner product B_V^+ on V by setting $\{e_1, \ldots, e_l, f_1, \ldots, f_l\}$ to be an orthonormal basis of $E \oplus F$, and $(E \oplus F)^\perp = U$.

If we set $K = G(V,B) \cap G(V, B_V^+)$, then K is a maximal compact subgroup of $G(V, B_V)$. Moreover, if we set

$$a(\lambda_1, \ldots, \lambda_l) = \begin{bmatrix} \lambda_1 & & & & & & \\ & \ddots & & & & & \\ & & \lambda_l & & & & \\ & & & I_{\dim U} & & & \\ & & & & \lambda_l^{-1} & & \\ & & & & & \ddots & \\ & & & & & & \lambda_1^{-1} \end{bmatrix},$$

then

$$A = \{a(\lambda_1, \ldots, \lambda_l) \mid \lambda_i \in \mathbb{R}^*, \text{ for } i = 1, \ldots, l\}$$

is a maximal split torus in $G(V, B_V)$ and, according to the Cartan decomposition, for any $g \in G(V, B_V)$, there exists k_1, $k_2 \in K$, and $a \in A$ such that $g = k_1 a k_2$.

Given $g \in G(V, B_V)$, let

$$\|g\| = \sup_{B_V^+(v,v)=1} [B_V^+(gv, gv)]^{\frac{1}{2}}$$

be the operator norm restricted to $G(V, B)$. Observe that if $g = k_1 a k_2$, with k_1, $k_2 \in K$ and $a = a(\lambda_1, ..., \lambda_l)$, then

$$\|g\| = \max\{|\lambda_1|, \dots, |\lambda_l|, |\lambda_1|^{-1}, \dots, |\lambda_l|^{-1}\}.$$

From this observation it is immediate that $\|g\| = \|g^{-1}\|$ and $\|\exp tX\| = \|\exp X\|^t$ for all $t \geq 0$, and all $X \in \mathfrak{a} = \mathrm{Lie}(A)$. Since $\|\cdot\|$ is the operator norm in $\mathrm{End}(V)$ we also have that $\|g_1 g_2\| \leq \|g_1\|\|g_2\|$ for all g_1, $g_2 \in G$, and hence $\|\cdot\|$ satisfies all the properties of a norm on G [41, 2.A.2].

4.2. **Inequalities regarding norms.** Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ be an $\mathfrak{sl}_2$-triple, and let $P = MN$ be as in Section 3.1. Assume that $V = \oplus_{k=-r}^{r} V_k$, and set $\mathrm{End}_i(V) = \oplus_{k=-r}^{r} \mathrm{Hom}(V_k, V_{k+i})$, where $V_k = 0$ if $|k| > r$. Then

$$\mathrm{End}(V) = \bigoplus_{i=-2r}^{2r} \mathrm{End}_i(V),$$

and $\mathfrak{g}_i = \mathfrak{g} \cap \mathrm{End}_i(V)$. Recall that, if $\mathfrak{g}_{-1} \neq 0$, then there is a symplectic structure on $\mathfrak{g}_{-1}$ given by $\kappa_{-1}(S, T) = \mathrm{Tr}(X[S, T])$. Let $\mathfrak{g}_{-1} = \mathfrak{e} \oplus \mathfrak{f}$ be a complete polarization of $\mathfrak{g}_{-1}$ with respect to this symplectic structure, and let $\mathrm{Her}_{-1}(V) = \{T \in \mathrm{End}_{-1}(V) \mid T^* = T\}$. (Here Her_{-1} stands for Hermitian of degree -1.) Then we have a decomposition $\mathrm{End}_{-1}(V) = \mathfrak{g}_{-1} \oplus \mathrm{Her}_{-1}(V) = \mathfrak{e} \oplus \mathfrak{f} \oplus \mathrm{Her}_{-1}(V)$. Define a norm $\|\cdot\|_{-1}$ on $\mathrm{End}_{-1}(V)$ by setting

$$\|T_{\mathfrak{e}} + T_{\mathfrak{f}} + T_{\mathrm{Her}_{-1}(V)}\|_{-1} = \|T_{\mathfrak{e}}\| + \|T_{\mathfrak{f}}\| + \|T_{\mathrm{Her}_{-1}(V)}\|,$$

where, as before, $\|\cdot\|$ is the operator norm, $T_{\mathfrak{e}} \in \mathfrak{e}$, $T_{\mathfrak{f}} \in \mathfrak{f}$, and $T_{\mathrm{Her}_{-1}(V)} \in \mathrm{Her}_{-1}(V)$ are arbitrary. Now given $T \in \mathrm{End}(V)$, let

$$\|T\|_\gamma = \|T_{-1}\|_{-1} + \sum_{i \neq -1} \|T_i\|,$$

where $T = \oplus T_i$, with $T_i \in \mathrm{End}_i(V)$. Then $\|\cdot\|_\gamma$ defines a norm on $\mathrm{End}(V)$ and, since all norms on a finite dimensional vector space are equivalent, there exists constants C_1, $C_2 > 0$ such that for all $T \in \mathrm{End}(V)$

$$C_1\|T\| \leq \|T\|_\gamma \leq C_2\|T\|. \tag{4.1}$$

Recall that $\mathfrak{n} = \mathfrak{u} \oplus \mathfrak{g}_{-1} = \mathfrak{u} \oplus \mathfrak{e} \oplus \mathfrak{f}$. Let $\mathfrak{n}_{\mathfrak{e}} = \mathfrak{u} \oplus \mathfrak{e}$ and $\mathfrak{n}_{\mathfrak{f}} = \mathfrak{u} \oplus \mathfrak{f}$, which are ideals of $\mathfrak{n}$. Let

$$N_{\mathfrak{e}} = \exp \mathfrak{n}_{\mathfrak{e}} \quad \text{and} \quad N_{\mathfrak{f}} = \exp \mathfrak{n}_{\mathfrak{f}} \tag{4.2}$$

be the corresponding normal subgroups of N. Observe that for every $n \in N$ there exists unique $u \in U$, $Z_{\mathfrak{f}} \in \mathfrak{f}$ and $Z_{\mathfrak{e}} \in \mathfrak{e}$ such that $n = u(\exp Z_{\mathfrak{f}})(\exp Z_{\mathfrak{e}})$. It follows immediately that $N_{\mathfrak{f}} \backslash N \cong \mathfrak{e}$.

Lemma 4.1. *For all $Z \in \mathfrak{e}$, $\tilde{n} \in N_{\mathfrak{f}}$, we have*

$$\|\tilde{n} \exp Z\| \geq C_1(1 + \|Z\|). \tag{4.3}$$

Proof. Observe that $\tilde{n} = \exp Z_1 \exp Z_2$, for some $Z_1 \in \mathfrak{u}$, $Z_2 \in \mathfrak{f}$. Hence, as an element of $\mathrm{End}(V)$, $\tilde{n} \exp Z = \exp Z_1 \exp Z_2 \exp Z = 1 + Z + Z_2 + \tilde{Z}$, for some $\tilde{Z} \in \oplus_{k=2}^{2r} \mathrm{End}_{-k}(V)$. Therefore,

$$\begin{aligned} \|\tilde{n} \exp Z\|_\gamma &= \|1 + Z + Z_1 + \tilde{Z}\|_\gamma \\ &= 1 + \|Z\| + \|Z_1\| + \|\tilde{Z}\|_\gamma \\ &\geq 1 + \|Z\|. \end{aligned}$$

The lemma then follows from (4.1). □

Lemma 4.2. *There exists constants $\tilde{d}$, $\tilde{C} > 0$ such that for all $Z \in \mathfrak{e}$, $\tilde{n} \in N_{\mathfrak{f}}$*

$$\|\tilde{n}(\exp Z)\|^{\tilde{d}} \geq \tilde{C}\|\tilde{n}\|\|(\exp Z)\|.$$

Proof. Since $\|\cdot\|$ is the operator norm on $\mathrm{End}(V)$, we have that

$$\|\exp Z\| \leq 1 + \|Z\| + \frac{\|Z\|^2}{2!} + \cdot + \frac{\|Z\|^{2r}}{2r!} \leq C_3(1+\|Z\|)^{2r}, \tag{4.4}$$

for some constant $C_3 > 0$. Here we have used that $V = \oplus_{k=-r}^{r} V_k$. Combining equations (4.3) and (4.4) we get that

$$\|\tilde{n}(\exp Z)\|^{2r} \geq C_1^{2r}(1+\|Z\|)^{2r} \geq \frac{C_1^{2r}}{C_3}\|\exp Z\|. \tag{4.5}$$

On the other hand, since the restriction of $\|\cdot\|$ to G defines a norm on G, we have that $\|\tilde{n}\| \leq \|(\exp Z)^{-1}\|\|\tilde{n}(\exp Z)\| = \|\exp Z\|\|\tilde{n}(\exp Z)\|$. From this and equation (4.5)

$$\|\tilde{n}(\exp Z)\| \geq \frac{\|\tilde{n}\|}{\|\exp Z\|} \geq \frac{C_1^{2r}}{C_3}\frac{\|\tilde{n}\|}{\|\tilde{n}(\exp Z)\|^{2r}},$$

and hence $\|\tilde{n}(\exp Z)\|^{2r+1} \geq \frac{C_1^{2r}}{C_3}\|\tilde{n}\|$. But now this inequality and equation (4.5) imply that, if we set $\tilde{C} = (\frac{C_1^{2r}}{C_3})^2$ and $\tilde{d} = 4r+1$, then

$$\|\tilde{n}(\exp Z)\|^{\tilde{d}} \geq \tilde{C}\|\tilde{n}\|\|\exp Z\|.$$

□

Corollary 4.3. *There exists constants* d_0, $C_0 > 0$ *such that, for all* $k \in K$, $m \in M$, $Z \in \mathfrak{e}$, *and* $\tilde{n} \in N_{\mathfrak{f}}$

$$\|\tilde{n}(\exp Z)mk\|^{d_0} \geq C_0\|\tilde{n}\|\|\exp Z\|\|m\|.$$

Proof. By definition of $\|\cdot\|$, $\|\tilde{n}(\exp Z)mk\| = \|\tilde{n}(\exp Z)m\|$. Now, from the proof of [41, Theorem 7.2.1], we have that

$$\|\tilde{n}(\exp Z)m\| \geq \|m\| \text{ and } \|\tilde{n}(\exp Z)m\|^2 \geq \|\tilde{n}(\exp Z)\|. \tag{4.6}$$

Let $\tilde{C}$ and $\tilde{d}$ be as in Lemma 4.2. Then, if we set $d_0 = 2\tilde{d}+1$ and $C_0 = \tilde{C}$, we have that

$$\|\tilde{n}(\exp Z)mk\|^{d_0} \geq \|\tilde{n}(\exp Z)\|^{\tilde{d}}\|m\| \geq C_0\|\tilde{n}\|\|\exp Z\|\|m\|.$$

□

4.3. **Some spaces of rapidly decreasing functions on G.** We retain all of the notation from the previous section.

Let $(\rho_{\chi_\gamma}, \mathscr{S}_{\chi_\gamma})$ be the smooth Heisenberg representation of N associated to the character χ_γ of U, as in (3.12). We first give a realization of $\mathscr{S}_{\chi_\gamma}$, as follows: fix $\mathfrak{g}_{-1} = \mathfrak{e} \oplus \mathfrak{f}$ (a complete polarization of $\mathfrak{g}_{-1}$). Set $\mathscr{D}(\mathfrak{e})$ to be the space of constant-coefficient differential operators on $\mathfrak{e}$, and let

$$\mathscr{S}(\mathfrak{e}) = \{f \in C^\infty(\mathfrak{e}) \mid p_{Z,d}(f) < \infty \text{ for all } Z \in \mathscr{D}(\mathfrak{e}),\, d \in \mathbb{N}\}$$

be the Schwartz space of $\mathfrak{e}$, where

$$p_{Z,d}(f) = \sup_{T \in \mathfrak{e}} |Zf(T)|(1+\|T\|)^d. \tag{4.7}$$

Extend χ_γ to $N_{\mathfrak{f}}$ by setting

$$\chi_\gamma(u \exp Z) = \chi_\gamma(u), \qquad \text{for all } u \in U,\, Z \in \mathfrak{f}.$$

Given $f \in \mathbb{C}^\infty(N_{\mathfrak{f}}\backslash N;\chi_\gamma)$, define $\hat{f} \in C^\infty(\mathfrak{e})$ by $\hat{f}(Z) = f(\exp Z)$, for all $Z \in \mathfrak{e}$. Conversely, given $f \in C^\infty(\mathfrak{e})$, set $\check{f}(n \exp Z) = \chi_\gamma(n)f(Z)$ for all $n \in N_{\mathfrak{f}}$, $Z \in \mathfrak{e}$. Clearly these two maps are inverse of each other, and if we set

$$\mathscr{S}(N_{\mathfrak{f}}\backslash N;\chi_\gamma) = \{f \in C^\infty(N_{\mathfrak{f}}\backslash N;\chi_\gamma) \mid \hat{f} \in \mathscr{S}(\mathfrak{e})\},$$

then

$$\mathscr{S}_{\chi_\gamma} \cong \mathscr{S}(N_{\mathfrak{f}}\backslash N;\chi_\gamma). \tag{4.8}$$

In the rest of this article we will frequently use this realization of $\mathscr{S}_{\chi_\gamma}$ (implicitly).

Remarks 4.4. (a) Here, and in the sequel, for a smooth representation σ of a closed subgroup B of a Lie group A,

$$C^\infty(B\backslash A;\sigma) = \{f \in C^\infty(A;\sigma) \,|\, f(ba) = \sigma(b)f(a), \text{for all } b \in B \text{ and } a \in A\}$$

denotes space of the representation of A smoothly induced from σ. Later, we will also need to consider the space $C_c^\infty(B\backslash A;\sigma)$ consisting of those elements in $C^\infty(B\backslash A;\sigma)$ with compact support modulo B.
(b) In the literature, the Lie algebras $\mathfrak{u}$, $\mathfrak{n}$ are frequently denoted by $\mathfrak{n}_2$ and $\mathfrak{n}_1$, respectively. When that is the case, the Lie algebra $\mathfrak{n}_{\mathfrak{f}}$ is often denoted $\mathfrak{n}_{1.5}$.
(c) When $\mathfrak{k}$ is non-Archimedian, we set $\mathscr{S}(\mathfrak{e})$ to be the space of all the locally constant, compactly supported functions on $\mathfrak{e}$ (known as the Bruhat-Schwartz space on $\mathfrak{e}$). With this definition equation (4.8) remains valid for $\mathfrak{k}$ non-Archimedian.

Let $U(\mathfrak{g})$ be the universal enveloping algebra of $\mathfrak{g}$. Given $f \in C^\infty(G)$, $Z \in U(\mathfrak{g})$ and $d \in \mathbb{N}$, we set

$$q_{Z,d}(f) = \sup_{g\in G} |R_Z f(g)| \|g\|^d,$$

where R_Z acts on $C^\infty(G)$ via the right regular representation of $U(\mathfrak{g})$. Let

$$\mathscr{C}(G) = \{f \in C^\infty(G) \mid q_{X,d}(f) < \infty, \text{ for all } X \in U(\mathfrak{g}),\, d \in \mathbb{N}\}\ .$$

It is easy to check that $\mathscr{C}(G)$ is a Fréchet space and that $q_{Z,d}$ is a semi-norm on $\mathscr{C}(G)$ for all $Z \in U(\mathfrak{g})$, $d \in \mathbb{N}$. We call $\mathscr{C}(G)$ the *space of rapidly decreasing functions* on G.

We now define certain space of rapidly decreasing functions on $N\backslash G$. Given $f \in C^\infty(N\backslash G;\mathscr{S}_{\chi_\gamma})$, $Z_1 \in U(\mathfrak{g})$, $Z_2 \in \mathscr{D}(\mathfrak{e})$, d_1, $d_2 \in \mathbb{N}$, set

$$q_{Z_1,Z_2,d_1,d_2}(f) = \sup_{k\in K,\, m\in M,\, T\in\mathfrak{e}} |Z_2(R_{Z_1}f(mk))(T)|(1+\|T\|)^{d_2}\|m\|^{d_1}. \tag{4.9}$$

Then we define

$$\mathscr{C}(N\backslash G;\mathscr{S}_{\chi_\gamma}) = \{f \in C^\infty(N\backslash G;\mathscr{S}_{\chi_\gamma}) \,|\, q_{Z_1,Z_2,d_1,d_2}(f) < \infty \text{ for all } q_{Z_1,Z_2,d_1,d_2} \text{ as in (4.9)}\}.$$

Observe that

$$q_{Z_1,Z_2,d_1,d_2}(f) = \sup_{k\in K, m\in M} p_{Z_2,d_2}(R_{Z_1}f(mk))\|m\|^{d_1},$$

where p_{Z_2,d_2} is as in (4.7). In general, given a smooth representation of N, $(\tau,\mathscr{H})$, and a seminorm ρ on $\mathscr{H}$, we set

$$q_{Z,d,\rho}(f) = \sup_{k\in K, m\in M} \rho(R_Z f(mk))\|m\|^d, \tag{4.10}$$

and define

$$\mathscr{C}(N\backslash G;\mathscr{H}) = \{f \in C^\infty(N\backslash G;\mathscr{H}) \,|\, q_{Z,d,\rho}(f) < \infty \text{ for all } q_{Z,d,\rho} \text{ as in (4.10)}\}.$$

Recall that there exists a covering $M_{\chi_\gamma} \twoheadrightarrow M_X$ such that $(\rho_{\chi_\gamma},\mathscr{S}_{\chi_\gamma})$ extends to a representation of $M_{\chi_\gamma} \ltimes N$. Using this extension, we define a natural action of M_{χ_γ} on $\mathscr{C}(N\backslash G;\mathscr{S}_{\chi_\gamma})$ by

$$m \cdot f(g) = \rho_{\chi_\gamma}(m) f(\bar{m}^{-1}g), \qquad \text{for all } m \in M_{\chi_\gamma},\, g \in G. \tag{4.11}$$

Here $\bar{m}$ is the image of m under the map $M_{\chi_\gamma} \twoheadrightarrow M_X$.

Now, given $f \in C^\infty(N_{\mathfrak{f}} \backslash G; \chi_\gamma)$, $Z \in U(\mathfrak{g})$ and $d \in \mathbb{N}$, we set

$$q_{Z,d}(f) = \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_Z f((\exp T)mk)| \|(\exp T)mk\|^d,$$

and define

$$\mathscr{C}(N_{\mathfrak{f}} \backslash G; \chi_\gamma) = \{f \in C^\infty(N_{\mathfrak{f}} \backslash G; \chi_\gamma) \,|\, q_{Z,d}(f) < \infty \text{ for all } Z \in U(\mathfrak{g}), d \in \mathbb{N}\}.$$

Lemma 4.5. *As G-modules, we have $\mathscr{C}(N_{\mathfrak{f}} \backslash G; \chi_\gamma) \cong \mathscr{C}(N \backslash G; \mathscr{S}_{\chi_\gamma})$.*

Proof. Given $f \in \mathscr{C}(N \backslash G; \mathscr{S}_{\chi_\gamma})$, define $\hat{f} \in C^\infty(N_{\mathfrak{f}} \backslash G; \chi_\gamma)$ by $\hat{f}(g) = f(g)(0)$. We claim that $\hat{f} \in \mathscr{C}(N_{\mathfrak{f}} \backslash G; \chi_\gamma)$. Effectively, given $Z \in U(\mathfrak{g})$, $d \in \mathbb{N}$,

$$\begin{aligned}
q_{Z,d}(\hat{f}) &= \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_Z \hat{f}((\exp T)mk)| \|(\exp T)mk\|^d \\
&= \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_Z f(nmk)(0)| \|(\exp T)mk\|^d \\
&\leq \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_Z f(mk)(T)| \|mk\|^d \|\exp T\|^d \\
&\leq \sup_{k \in K, m \in M, T \in \mathfrak{e}} C_3 |R_Z f(mk)(T)| \|mk\|^d (1 + \|T\|)^{2rd} \\
&\leq C_3 q_{Z,1,d,2rd}(f) < \infty,
\end{aligned}$$

where we have used equation (4.4) in the next to last equality.

Now given $f \in \mathscr{C}(N_{\mathfrak{f}} \backslash G; \chi_\gamma)$, set $\check{f}(g)(T) = f(\exp(T)\, g)$, for all $g \in G$, $T \in \mathfrak{e}$. We claim that $\check{f} \in \mathscr{C}(N \backslash G; \mathscr{S}_{\chi_\gamma})$. Effectively, given $Z_1 \in U(\mathfrak{g})$, $Z_2 \in \mathscr{D}(\mathfrak{e})$, d_1, $d_2 \in \mathbb{N}$, we have that

$$\begin{aligned}
q_{Z_1,Z_2,d_1,d_2}(\check{f}) &= \sup_{k \in K, m \in M, T \in \mathfrak{e}} |Z_2(R_{Z_1} \check{f}(mk))(T)| \|mk\|^{d_1} (1 + \|T\|)^{d_2} \\
&= \sup_{k \in K, m \in M, T \in \mathfrak{e}} |(R_{\mathrm{Ad}(mk)^{-1} Z_2 Z_1} f(\exp(T)mk)| \|mk\|^{d_1} (1 + \|T\|)^{d_2},
\end{aligned}$$

where we have identified Z_2 with an element in $U^l(\mathfrak{g})$ for some l. Let $\{\tilde{Z}_1, \ldots, \tilde{Z}_s\}$ be a basis of $U^l(\mathfrak{g})$. Then

$$\mathrm{Ad}(mk)^{-1} Z_2 = \sum_{j=1}^{s} a_j(mk) \tilde{Z}_j,$$

for some functions a_j. Since $(\mathrm{Ad}, U^l(\mathfrak{g}))$ is finite dimensional, it is of moderate growth, and hence there exists constants C_l, $d_l > 0$ such that $|a_j(mk)| \leq C_l \|mk\|^{d_l}$, for all $k \in K$, $m \in M$. From this, equations (4.3) and (4.6), we have

$$\begin{aligned}
q_{Z_1,Z_2,d_1,d_2}(\check{f}) &\leq C_l \sum_{j=1}^{s} \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_{\tilde{Z}_j Z_1} f(\exp(T)mk)| \|mk\|^{d_1 + d_l} (1 + \|T\|)^{d_2} \\
&\leq \frac{C_l}{C_1^{d_2}} \sum_{j=1}^{s} \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_{\tilde{Z}_j Z_1} f(\exp(T)mk)| \|mk\|^{d_1 + d_l} \|\exp T\|^{d_2} \\
&\leq \frac{C_l}{C_1^{d_2}} \sum_{j=1}^{s} \sup_{k \in K, m \in M, T \in \mathfrak{e}} |R_{\tilde{Z}_j Z_1} f(\exp(T)mk)| \|\exp(T)mk\|^{d_1 + d_l + 2d_2} \\
&\leq \frac{C_l}{C_1^{d_2}} \sum_{j=1}^{s} q_{\tilde{Z}_j Z_1, d_1 + d_l + 2d_2}(f) < \infty.
\end{aligned}$$

Finally, we easily check that $\check{\hat{f}} = f$ and $\hat{\check{f}} = f$. □

Given $f \in \mathscr{C}(G)$, define a function $f_{\chi_\gamma} \in C^\infty(N_{\mathfrak{f}}\backslash G;\chi_\gamma)$ by

$$f_{\chi_\gamma}(g) = \int_{N_{\mathfrak{f}}} f(\tilde{n}g)\chi_\gamma(\tilde{n})^{-1}\,d\tilde{n}.$$

Lemma 4.6. *The map $f \mapsto f_{\chi_\gamma}$ defines a surjective G-intertwining map from $\mathscr{C}(G)$ to the space $\mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_\gamma)$.*

Proof. We will first show that $f_{\chi_\gamma} \in \mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_\gamma)$. Given $Z \in U(\mathfrak{g})$, $d \in \mathbb{N}$, we have that

$$\begin{aligned} q_{Z,d}(f_{\chi_\gamma}) &= \sup_{k\in K, m\in M, T\in\mathfrak{e}} |R_Z f_{\chi_\gamma}((\exp T)mk)|\|(\exp T)mk\|^d \\ &\leq \sup_{k,m,T} \int_{N_{\mathfrak{f}}} |R_Z f_{\chi_\gamma}(\tilde{n}(\exp Z)mk)|\,d\tilde{n}\,\|(\exp T)mk\|^d. \end{aligned}$$

Now we know that for all $d_1 \in \mathbb{N}$, $|R_Z f(\tilde{n}(\exp Z)mk)|\|\tilde{n}(\exp Z)mk\|^{d_1} \leq q_{Z,d_1}(f)$. Hence, from Corollary 4.3,

$$\begin{aligned} q_{Z,d}(f_{\chi_\gamma}) &\leq \sup_{k\in K, m\in M, T\in\mathfrak{e}} q_{Z,d_1}(f)\int_{N_{\mathfrak{f}}} \|\tilde{n}(\exp Z)mk\|^{-d_1}\,d\tilde{n}\|(\exp T)mk\|^d \\ &\leq \sup_{k\in K, m\in M, T\in\mathfrak{e}} q_{Z,d_1}(f)\int_{N_{\mathfrak{f}}} (C_0\|\tilde{n}\|\|\exp T\|\|m\|)^{-\frac{d_1}{d_0}}(\|\exp T\|\|m\|)^d\,d\tilde{n} \\ &= \sup_{k\in K, m\in M, T\in\mathfrak{e}} q_{Z,d_1}(f)C_0^{-\frac{d_1}{d_0}}(\|\exp T\|\|m\|)^{d-\frac{d_1}{d_0}}\int_{N_{\mathfrak{f}}} \|\tilde{n}\|^{-\frac{d_1}{d_0}}\,d\tilde{n}. \end{aligned}$$

But it is clear that, if $d_1 \gg 0$, then the right hand side of the above equation is finite.

Now we will show that the map is surjective. Fix a function $\phi \in C_c^\infty(N_{\mathfrak{f}})$ such that

$$\int_{N_{\mathfrak{f}}} \phi(\tilde{n})\chi_\gamma(\tilde{n})^{-1}\,d\tilde{n} = 1.$$

Given $f \in \mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_\gamma)$, let $h \in C^\infty(G)$ be given by $h(\tilde{n}(\exp Z)mk) = \phi(\tilde{n})f(nmk)$. Then it is clear that $h \in \mathscr{C}(G)$ and $h_{\chi_\gamma}(nmk) = f(nmk)$ for all $k \in K$, $m \in M$ and $Z \in \mathfrak{e}$. From all this we conclude that the map $f \mapsto f_{\chi_\gamma}$ is surjective. ☐

Remark 4.7. When $\mathfrak{k}$ is non-Archimedian, we set $\mathscr{C}(G)$ to be the space of locally constant, compactly supported functions, with analogous definitions for $\mathscr{C}(N\backslash G;\mathscr{S}_{\chi_\gamma})$ and $\mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_\gamma)$. With these definitions, it is straightforward to check that all the results in this section remain valid in the non-Archimedian case.

4.4. **Realizing a generalized Whittaker model on $N\backslash G$.** In this section, we give the promised realization of the space of generalized Whittaker models for a Casselman-Wallach representation of G. But before stating this result, we need the following technical lemma.

Lemma 4.8. *Let $(\pi,\mathscr{V})$ be a Casselman-Wallach representation of G. Then*

$$(\mathscr{V}')^{N_{\mathfrak{f}},\chi_\gamma} \cong \mathrm{Wh}_\gamma(\pi).$$

Proof. Given $\lambda \in \mathrm{Wh}_\gamma(\pi)$, set $\hat{\lambda}(v) = \lambda(v)(0)$, for $v \in \mathscr{V}$. Then it is clear that $\hat{\lambda} \in (\mathscr{V}')^{N_{\mathfrak{f}},\chi_\gamma}$. On the other hand, given $\lambda \in (\mathscr{V}')^{N_{\mathfrak{f}},\chi_\gamma}$, set $\check{\lambda}(v)(T) = \lambda(\pi(\exp T)\,v)$ for $v \in \mathscr{V}$, $T \in \mathfrak{e}$. Then it is clear that $\check{\lambda}(v) \in C^\infty(\mathfrak{e})$, but we claim that it is actually in $\mathscr{S}(\mathfrak{e})$. To see this, observe that if $R \in \mathfrak{e}$, then

$$\check{\lambda}(d\pi(R)v)(T) = D_R\check{\lambda}(v)(T), \qquad \text{for all } T \in \mathfrak{e}, \tag{4.12}$$

where $D_R \in \mathscr{D}(\mathfrak{e})$ represents the derivative in the direction R. On the other hand, if $S \in \mathfrak{f}$, then

$$\check{\lambda}(d\pi(S)v)(T) = d\chi_\gamma(\kappa(S,T))\check{\lambda}(v)(T), \qquad \text{for all } T \in \mathfrak{e}, \tag{4.13}$$

where $d\chi_\gamma$ is the linear functional given by the derivative of the character χ_γ. But now, since $(\pi, \mathscr{V})$ is a representation of moderate growth, we can find a constant $d > 0$ such that for all $v \in \mathscr{V}$, there exists $C_{\lambda,v} > 0$ such that

$$|\check{\lambda}(v)(T)| = |\lambda(\pi(\exp T)\, v)| \leq C_{\lambda,v} \|\exp T\|^d \leq C_3 C_{\lambda,v} (1 + \|T\|)^{2dr}, \qquad \text{for all } T \in \mathfrak{e}. \tag{4.14}$$

Here C_3 and r are as in equation (4.4). Observe that, although the constant $C_{\lambda,v}$ depends on v (and on λ), d is independent of the element $v \in \mathscr{V}$ chosen. Therefore, from equations (4.12), (4.13) and (4.14) for all $v \in \mathscr{V}$ the function $\check{\lambda}(v)$ is such that if we take any partial derivatives, and multiply by any polynomial on $\mathfrak{e}$, it still has growth controlled by a polynomial of degree $2dr$. But this implies that $\check{\lambda}(v) \in \mathscr{S}(\mathfrak{e})$, as we wanted to show.

To finish the proof we just have to show that $\hat{\check{\lambda}} = \lambda$, and $\check{\hat{\lambda}} = \lambda$, but this follow easily from the definitions. ☐

Given an $\mathfrak{sl}_2$-triple $\gamma = \{X, H, Y\} \subset \mathfrak{g}$, set

$$\check{\gamma} = \{-X, H, -Y\}. \tag{4.15}$$

Then, it is immediate to check that $\check{\gamma}$ is also an $\mathfrak{sl}_2$-triple and that $\chi_{\check{\gamma}} = \check{\chi}_\gamma$, where $\check{\chi}_\gamma$ is the character of U given by $\check{\chi}_\gamma(u) = \chi_\gamma(u)^{-1}$ for all $u \in U$.

Proposition 4.9. *Let $(\pi, \mathscr{V})$ be an irreducible Casselman-Wallach representation of G, and let $\mathscr{C}(N\backslash G; \mathscr{S}_{\chi_{\check{\gamma}}})_{G,\pi}$ denote the maximal π-isotypic quotient of $\mathscr{C}(N\backslash G; \mathscr{S}_{\chi_{\check{\gamma}}})$. Then, as an $M_{\chi_\gamma} \times G$-module,*

$$\mathscr{C}(N\backslash G; \mathscr{S}_{\chi_{\check{\gamma}}})_{G,\pi} \cong W_\gamma(\check{\pi}) \otimes \pi,$$

where $\check{\pi}$ is the contragredient Casselman-Wallach representation of π and $W_\gamma(\check{\pi})$ is the continuous dual of $\mathrm{Wh}_\gamma(\check{\pi})$. Here $\otimes$ stands for the completed projective tensor product (of two locally convex topological spaces).

Proof. By Lemma 4.5, we have $\mathscr{C}(N\backslash G; \mathscr{S}_{\chi_{\check{\gamma}}}) \cong \mathscr{C}(N_{\mathfrak{f}}\backslash G; \chi_{\check{\gamma}})$. Let $\lambda \in \mathrm{Hom}_G(\mathscr{C}(N_{\mathfrak{f}}\backslash G; \chi_{\check{\gamma}}), \mathscr{V})$, and let $(\check{\pi}, \check{\mathscr{V}})$ be the representation contragredient to $(\pi, \mathscr{V})$. Then we may think of λ as an element in a space of $\check{\mathscr{V}}'$-valued distributions. Hence, according to [14] there exists $\mu_\lambda \in (\check{\mathscr{V}}')^{N_{\mathfrak{f}}, \chi_\gamma}$ such that if $f \in C_c^\infty(N_{\mathfrak{f}}\backslash G; \chi_\gamma)$, then

$$\lambda(f)(v) = \int_{N_{\mathfrak{f}}\backslash G} f(g) \mu_\lambda(\pi(g)v)\, dg, \qquad \text{for } v \in \check{\mathscr{V}}.$$

On the other hand, given $\mu \in (\check{\mathscr{V}}')^{N_{\mathfrak{f}}, \chi_\gamma}$ and $f \in \mathscr{C}(N_{\mathfrak{f}}\backslash G; \chi_{\check{\gamma}})$, there exists $h \in \mathscr{C}(G)$ such that $f = h_{\chi_{\check{\gamma}}}$ and for all $v \in \check{\mathscr{V}}$

$$\begin{aligned}
\int_G h(g)\mu_\lambda(\pi(g)v)\, dg &= \int_{N_{\mathfrak{f}}\backslash G} \int_{N_{\mathfrak{f}}} h(\tilde{n}g)\mu_\lambda(\pi(\tilde{n})\pi(g)v)\, d\tilde{n}\, dg \\
&= \int_{N_{\mathfrak{f}}\backslash G} \int_{N_{\mathfrak{f}}} h(\tilde{n}g)\chi_\gamma(\tilde{n})\, d\tilde{n}\ \mu_\lambda(\pi(g)v)\, dg \\
&= \int_{N_{\mathfrak{f}}\backslash G} \int_{N_{\mathfrak{f}}} h(\tilde{n}g)\chi_{\check{\gamma}}(\tilde{n})^{-1}\, d\tilde{n}\ \mu_\lambda(\pi(g)v)\, dg \\
&= \int_{N_{\mathfrak{f}}\backslash G} h_{\chi_{\check{\gamma}}}(g)\mu_\lambda(\pi(g)v)\, dg \\
&= \int_{N_{\mathfrak{f}}\backslash G} f(g)\mu_\lambda(\pi(g)v)\, dg.
\end{aligned}$$

Therefore, if we set

$$\lambda_\mu(f)(v) = \int_{N_{\mathfrak{f}}\backslash G} f(g)\mu_\lambda(\pi(g)v)\,dg, \qquad \text{for } v \in \check{\mathscr{V}},$$

then $\lambda_\mu \in \operatorname{Hom}_G(\mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_{\check{\gamma}}), \check{\mathscr{V}}')$. Method of [36, Section 3] implies that $\lambda_\mu(f)$ extends to a continuous linear functional on $\mathscr{V}'$, i.e., $\lambda_\mu(f)$ is actually an element of $\mathscr{V}$ for all $f \in \mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_{\check{\gamma}})$, or equivalently $\lambda_\mu \in \operatorname{Hom}_G(\mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_{\check{\gamma}}), \mathscr{V})$. Now it is easy to check that $\lambda_{\mu_\lambda} = \lambda$ and $\mu_{\lambda_\mu} = \mu$. Therefore

$$\mathscr{C}(N_{\mathfrak{f}}\backslash G;\chi_{\check{\gamma}})_{G,\pi} \cong \pi \otimes \check{\pi}_{N_{\mathfrak{f}},\chi_\gamma}.$$

Finally, according to Lemma 4.8, $\check{\pi}_{N_{\mathfrak{f}},\chi_\gamma} \cong W_\gamma(\check{\pi})$ and the proposition follows. □

Remarks 4.10. (a) In the Archimedian setting whenever we take the tensor product of two locally convex topological vector spaces, we always mean the completed projective tensor product. Observe that if $(\pi, \mathscr{V})$ is a Casselman-Wallach representation, then $\mathscr{V}$ is nuclear, and hence the completed projective tensor product of $\mathscr{V}$ with any locally convex topological vector space is equivalent to the completed injective tensor product [5].
(b) Proposition 4.9 also holds for $\Bbbk$ non-Archimedian, if we replace in its statement Casselman-Wallach representations by smooth, finitely generated, admissible representations, and completed projective tensor product by algebraic tensor product. The proof for this case follows the same line, but it is more straightforward.

5. Dual pairs and theta lifting of nilpotent orbits

5.1. **Reductive dual pairs of type I.** Let (V, B) be an ε-Hermitian module, and let $(\tilde{V}, \tilde{B})$ be an $\tilde{\varepsilon}$-Hermitian module such that $\varepsilon\tilde{\varepsilon} = -1$. Let $G = G(V)$ and $\tilde{G} = G(\tilde{V})$ be its associated isometry groups. We will use the ε-Hermitian form B to identify the $\Bbbk$-vector spaces $\tilde{V} \otimes_D V^*$ and $\operatorname{Hom}_D(V, \tilde{V})$. Given $T \in \operatorname{Hom}_D(V, \tilde{V})$, define $T^* \in \operatorname{Hom}_D(\tilde{V}, V)$ by

$$\tilde{B}(Tv, \tilde{v}) = B(v, T^*\tilde{v}) \qquad \text{for all } v \in V,\ \tilde{v} \in \tilde{V}.$$

Define similarly the $*$ map from $\operatorname{Hom}_D(\tilde{V}, V)$ to $\operatorname{Hom}_D(V, \tilde{V})$. Note that since $\varepsilon\tilde{\varepsilon} = -1$, we have $T^{**} = -T$ for $T \in \operatorname{Hom}_D(V, \tilde{V})$.

We now define a symplectic form $\langle \cdot, \cdot \rangle$ on $\operatorname{Hom}_D(V, \tilde{V})$ by setting

$$\langle T, S \rangle = \operatorname{Tr}(T^*S) \qquad \text{for all } T,\ S \in \operatorname{Hom}_D(V, \tilde{V}), \tag{5.1}$$

where $\operatorname{Tr}(T^*S)$ is the trace of T^*S as a $\Bbbk$-linear transformation. Let

$$\operatorname{Sp}(\operatorname{Hom}_D(V, \tilde{V})) = \left\{ g \in \operatorname{GL}(\operatorname{Hom}_D(V, \tilde{V}), \Bbbk) \;\middle|\; \begin{array}{c} \langle g \cdot T, g \cdot S \rangle = \langle T, S \rangle \\ \text{for all } T,\ S \in \operatorname{Hom}_D(V, \tilde{V}) \end{array} \right\}.$$

Then there is a natural map $G \times \tilde{G} \longrightarrow \operatorname{Sp}(\operatorname{Hom}_D(V, \tilde{V}))$ given by

$$(g, \tilde{g}) \cdot T = \tilde{g} T g^{-1} \qquad \text{for all } T \in \operatorname{Hom}(V, \tilde{V}),\ g \in G,\ \tilde{g} \in \tilde{G}.$$

We will use this map to identify G and $\tilde{G}$ with subgroups of $\operatorname{Sp}(\operatorname{Hom}_D(V, \tilde{V}))$. These two subgroups are mutual centralizers of each other, and form an example of a *reductive dual pair* of type I. See [7].

5.2. **Moment maps and lifting of nilpotent orbits.** Given $T \in \mathrm{Hom}_D(V, \tilde{V})$, it is clear that $T^*T \in \mathfrak{g}$ and $TT^* \in \tilde{\mathfrak{g}}$. Following [16, 3] we define the *moment maps* to be

$$\begin{array}{rcl} \varphi : \mathrm{Hom}_D(V, \tilde{V}) & \longrightarrow & \mathfrak{g} \\ T & \mapsto & T^*T \end{array}$$

and

$$\begin{array}{rcl} \tilde{\varphi} : \mathrm{Hom}_D(V, \tilde{V}) & \longrightarrow & \tilde{\mathfrak{g}} \\ T & \mapsto & TT^*. \end{array}$$

It is also clear that $\varphi(T)$ is nilpotent if and only if $\tilde{\varphi}(T)$ is nilpotent.

As in the introduction, let $\mathrm{Max}\,\mathrm{Hom}(V, \tilde{V})$ be the set of full rank elements in $\mathrm{Hom}(V, \tilde{V})$. Without any loss of generality, we assume that $\dim V \leq \dim \tilde{V}$, and elements of $\mathrm{Max}\,\mathrm{Hom}(V, \tilde{V})$ are then represented by injective maps from V to $\tilde{V}$. Recall also our standing assumption on the nilpotent orbit $\mathcal{O} \subset \mathfrak{g}$:

$$\varphi^{-1}(\mathcal{O}) \cap \mathrm{Max}\,\mathrm{Hom}(V, \tilde{V}) \neq \emptyset. \tag{5.2}$$

We first prove a result on dual pairs in the stable range, which says that in this case any (not just nilpotent) orbit satisfies the stated assumption. Recall that the dual pair $(G, \tilde{G})$ is in the stable range, with G the smaller member, if there is a polarization $\tilde{V} = \tilde{E} \oplus \tilde{U} \oplus \tilde{F}$, where $\tilde{E}$, $\tilde{F}$ are totally isotropic complementary subspaces with $\dim \tilde{E} = \dim \tilde{F} = \dim V$.

Lemma 5.1. *Assume that the dual pair $(G, \tilde{G})$ is in the stable range with G the smaller member. Given $X \in \mathfrak{g}$, there exists an injective map $T \in \mathrm{Hom}(V, \tilde{V})$ such that $T^*T = X$.*

Proof. Fix a linear isomorphism $T_{\tilde{E}} : V \to \tilde{E}$ and define a linear map $T_{\tilde{F}} : V \to \tilde{F}$ by $T_{\tilde{F}} = (T_{\tilde{E}}^*)^{-1}X/2$. Then, if we set $T = T_{\tilde{E}} + T_{\tilde{F}}$, we have that

$$T^*T = (T_{\tilde{E}}^* + T_{\tilde{F}}^*)(T_{\tilde{E}} + T_{\tilde{F}}) = T_{\tilde{E}}^*T_{\tilde{F}} + T_{\tilde{F}}^*T_{\tilde{E}} = X/2 + X/2 = X.$$

Note that we have used the fact that $T_{\tilde{F}}^* = X^*(T_{\tilde{E}}^{**})^{-1}/2 = XT_{\tilde{E}}^{-1}/2$. Now, since $T_{\tilde{E}}$ is injective, we conclude that T is injective and $T^*T = X$ as we wanted to show. □

Remark 5.2. The correspondence of nilpotent orbits for dual pairs in the stable range is well understood. See [3, 27], and [24] (for $K_{\mathbb{C}}$-nilpotent orbits). This is also covered by the next proposition, in view of Lemma 5.1.

Proposition 5.3. *Let $\mathcal{O} \subset \mathfrak{g}$ be a nilpotent orbit corresponding to the admissible ε-Hermitian Young tableau $\Gamma = (d^\Gamma, (V^\Gamma, B^\Gamma))$, where $d^\Gamma = [t_1^{i_1}, \ldots, t_l^{i_l}]$. Given $X \in \mathcal{O}$, let $T \in \mathrm{Hom}(V, \tilde{V})$ be an injective map such that $T^*T = X$. Then the orbit of the nilpotent element $\tilde{X} := TT^* \in \tilde{\mathfrak{g}}$ corresponds to the unique equivalence class of admissible $\tilde{\varepsilon}$-Hermitian Young tableaux $\tilde{\Gamma} = (d^{\tilde{\Gamma}}, (V^{\tilde{\Gamma}}, B^{\tilde{\Gamma}}))$, where*

- *$d^{\tilde{\Gamma}} = [(t_1+1)^{i_1}, \ldots, (t_l+1)^{i_l}, 1^s]$, with $s = \dim \tilde{V} - \dim V - \sum_{j=1}^{l} i_j$, namely $\tilde{\Gamma}$ is obtained by adding a column of length $\dim \tilde{V} - \dim V$ to Γ;*
- *$(V_{t_j}^{\tilde{\Gamma}, t_j+1}, B_{t_j}^{\tilde{\Gamma}, t_j+1}) \cong (V_{t_j-1}^{\Gamma, t_j}, B_{t_j-1}^{\Gamma, t_j})$, for all $j = 1, \ldots, l$;*
- *$(V_0^{\tilde{\Gamma},1}, B_0^{\tilde{\Gamma},1})$ is an $\tilde{\varepsilon}$-Hermitian module of dimension s.*

Proof. Let $\gamma = \{X, H, Y\}$ be an $\mathfrak{sl}_2$-triple containing X. Define V_i^{γ, t_j} as in Section 3.1, and set

$$\tilde{V}_{i+1}^{\tilde{\gamma}, t_j+1} = T(V_i^{\gamma, t_j}).$$

(At this point, $\tilde{\gamma}$ has no meaning.) Since T is injective, we have a direct sum decomposition:

$$T(V) = \bigoplus_{j=1}^{l} T(V^{\gamma,t_j}),$$

and for each j, we have

$$T(V^{\gamma,t_j}) = \bigoplus_{i=1}^{t_j} T(V^{\gamma,t_j}_{-t_j+2i-1}) = \bigoplus_{i=1}^{t_j} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i}.$$

Observe that for all v, $w \in V$,

$$\tilde{B}(Tv, Tw) = B(v, T^*Tw) = B(v, Xw).$$

This implies that $T(V^{\gamma,t_i}) \perp T(V^{\gamma,t_j})$ for $i \neq j$. We also note the following: if $\mathcal{A}, \mathcal{B} \subset V$ and $\mathcal{A}, X(\mathcal{B})$ form a perfect pairing under B, then $T(\mathcal{A}), T(\mathcal{B})$ form a perfect pairing under $\tilde{B}$. This implies that, for a fixed j, $\bigoplus_{i=1}^{t_j-1} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i}$ is non-degenerate, and $\tilde{V}^{\tilde{\gamma},t_j+1}_{t_j}$ is orthogonal to $\bigoplus_{i=1}^{t_j-1} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i}$. Therefore there exists $\tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j}$ such that

- $T(V^{\gamma,t_j}) \oplus \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j} = \bigoplus_{i=0}^{t_j} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i}$ is non-degenerate; and
- $\tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j} \perp T(V^{\gamma,t_i})$, for all $i \neq j$.

By an inductive argument, we may thus find $\tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j}$ for $1 \leq j \leq l$ such that

- $\tilde{U}_j =: \oplus_{i=0}^{t_j} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i}$ is non-degenerate; and
- $\oplus_{j=1}^{l} \tilde{U}_j$ is an orthogonal direct sum.

Let $\tilde{V}^{\tilde{\gamma},1}_0$ be the orthogonal complement of $\oplus_{j=1}^{l} \tilde{U}_j$. Then, we have a decomposition

$$\tilde{V} = \boxplus_{j=1}^{l} \left(\oplus_{i=0}^{t_j} \tilde{V}^{\tilde{\gamma},t_j+1}_{-t_j+2i} \right) \boxplus \tilde{V}^{\tilde{\gamma},1}_0,$$

where $\boxplus$ denotes an orthogonal direct sum.

Let $\tilde{V}_i = \oplus_j \tilde{V}^{\tilde{\gamma},t_j+1}_i$, and define an element $\tilde{H} \in \tilde{\mathfrak{g}}$ (easily checked) by setting $\tilde{H}\tilde{v} = i\,\tilde{v}$ for all $\tilde{v} \in \tilde{V}_i$. Note that for $i < t_j - 1$, $T^*|_{\tilde{V}^{\tilde{\gamma},t_j+1}_i} : \tilde{V}^{\tilde{\gamma},t_j+1}_i \to V^{\gamma,t_j}_{i+1}$ is an isomorphism and $TT^* = \tilde{X} : \tilde{V}^{\tilde{\gamma},t_j+1}_i \to \tilde{V}^{\tilde{\gamma},t_j+1}_{i+2}$. In particular, $\tilde{X}$ is nilpotent. Since $[\tilde{X}, \tilde{H}] = 2\tilde{X}$ and $\tilde{H}$ is semisimple, there exists an element $\tilde{Y} \in \tilde{\mathfrak{g}}$ such that $\{\tilde{X}, \tilde{H}, \tilde{Y}\}$ is an $\mathfrak{sl}_2$-triple. See [2, Section 3.3]. From all this it is clear that if $\tilde{\Gamma} = (d^{\tilde{\Gamma}}, (V^{\tilde{\Gamma}}, B^{\tilde{\Gamma}}))$ is the $\tilde{\varepsilon}$-Hermitian Young tableau associated to the orbit of $\tilde{X}$, then $d^{\tilde{\Gamma}}$ is obtained by adding a column to d^{Γ}. Finally observe that $\tilde{X}^{t_j} = TX^{t_j-1}T^*$, and so

$$\tilde{X}^{-t_j}|_{\tilde{V}^{\tilde{\gamma},t_j+1}_{t_j}} = (T^{-1})^* X^{-(t_j-1)} T^{-1}|_{\tilde{V}^{\tilde{\gamma},t_j+1}_{t_j}}.$$

If $\tilde{v}$, $\tilde{w} \in \tilde{V}^{\tilde{\gamma},t_j+1}_{t_j}$, then

$$\begin{aligned} \tilde{B}^{\tilde{\gamma},t_j+1}_{t_j}(\tilde{v}, \tilde{w}) &= \tilde{B}(\tilde{X}^{-t_j}\tilde{v}, \tilde{w}) \\ &= \tilde{B}((T^{-1})^* X^{-(t_j-1)} T^{-1}\tilde{v}, \tilde{w}) \\ &= B(X^{-(t_j-1)} T^{-1}\tilde{v}, T^{-1}\tilde{w}) \\ &= B^{\gamma,t_j}_{t_j+1}(T^{-1}\tilde{v}, T^{-1}\tilde{w}), \end{aligned} \tag{5.3}$$

which implies that $(V^{\tilde{\Gamma},t_j+1}_{t_j}, B^{\tilde{\Gamma},t_j+1}_{t_j}) \cong (V^{\Gamma,t_j}_{t_j-1}, B^{\Gamma,t_j}_{t_j-1})$, for all $j = 1, \ldots, l$. □

Let $\mathcal{O} \subset \mathfrak{g}$ be a nilpotent orbit (satisfying (5.2)), which corresponds to an ε-Hermitian Young tableau $\Gamma = (d^{\Gamma}, (V^{\Gamma}, B^{\Gamma}))$. We define the *theta lift* of $\mathcal{O}$, $\Theta(\mathcal{O}) \subset \tilde{\mathfrak{g}}$, to be the nilpotent orbit corresponding to the $\tilde{\varepsilon}$-Hermitian Young tableau $\tilde{\Gamma} = (d^{\tilde{\Gamma}}, (V^{\tilde{\Gamma}}, B^{\tilde{\Gamma}}))$ specified in Proposition 5.3.

Remark 5.4. It can be shown that, for the orbit $\mathcal{O}$ under consideration, $\tilde{\varphi}(\varphi^{-1}(\overline{\mathcal{O}})) = \overline{\Theta(\mathcal{O})}$. Cf. [16, Lemma 4.3]. Therefore, our definition of the theta lift of $\mathcal{O}$ agrees with the usual one.

5.3. **Lifting of $\mathfrak{sl}_2$-triples.** In this section we will introduce the concept of lifting of $\mathfrak{sl}_2$-triples. We start with the following

Definition 5.5. Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\tilde{\mathcal{O}}$, respectively. We say that $T \in \operatorname{Hom}(V, \tilde{V})$ *lifts* γ to $\tilde{\gamma}$ if $T^*T = X$, $TT^* = \tilde{X}$ and $T(V_j) \subset \tilde{V}_{j+1}$ for all j.

Set

$$\mathcal{O}_{\gamma,\tilde{\gamma}} = \{T \in \operatorname{Hom}(V, \tilde{V}) \,|\, T \text{ lifts } \gamma \text{ to } \tilde{\gamma}\}. \tag{5.4}$$

We prove the following elementary

Lemma 5.6. *Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples. Let $T \in \mathcal{O}_{\gamma,\tilde{\gamma}}$ and denote*

$$T_j = T|_{V_j} \in \operatorname{Hom}(V_j, \tilde{V}_{j+1}).$$

Then T_j is injective for $j < 0$, and surjective for $j \geq 0$.

Proof. We have $X|_{V_j} = T^*T|_{V_j} = T^*_{-j-2}T_j$. Since all highest weight vectors of a finite dimensional $\mathfrak{sl}_2$ representation have non-negative weights, we know $X|_{V_j}$ is injective for $j < 0$. This implies the first assertion.

For the second assertion, we first note that $T_j^* : \tilde{V}_{-j-1} \mapsto V_{-j}$. We have $\tilde{X}|_{\tilde{V}_{-j-1}} = TT^*|_{\tilde{V}_{-j-1}} = T_{-j}T_j^*$. By the same reasoning, we know that T_j^* is injective for $j \geq 0$. The second assertion follows. ☐

From now on, we assume that $\mathcal{O}$ satisfies (5.2), as in the introduction. We let

$$\mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}} = \mathcal{O}_{\gamma,\tilde{\gamma}} \cap \operatorname{Max}\operatorname{Hom}(V, \tilde{V}). \tag{5.5}$$

Lemma 5.7. *Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. Then $\mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}} \neq \emptyset$ and it is a single $M_X \times \tilde{M}_{\tilde{X}}$-orbit.*

Proof. From the proof of Proposition 5.3, we know $\mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}} \neq \emptyset$. Now let $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$ and set

$$T_i^{\gamma,t_j} = T|_{V_i^{\gamma,t_j}} \qquad \text{for all } 1 \leq j \leq l,\ -t_j + 1 \leq i \leq t_j - 1.$$

We claim that T_i^{γ,t_j} defines a linear isomorphism between V_i^{γ,t_j} and $\tilde{V}_{i+1}^{\tilde{\gamma},t_j+1}$ for all $1 \leq j \leq l$ and $-t_j + 1 \leq i \leq t_j - 1$. Since $TX = \tilde{X}T$, $T(V_i) \subset \tilde{V}_{i+1}$ and T is injective, we clearly have $T(V_{t_j-1}^{\gamma,t_j}) \subseteq \tilde{V}_{t_j}^{\tilde{\gamma},t_j+1}$. By Lemma 5.6, T_j is surjective for all $j \geq 0$. Thus any element of $\tilde{V}_{t_j}^{\tilde{\gamma},t_j+1}$ is of the form $T(v)$ for some $v \in V_{t_j-1}$. But then $\tilde{X}T(v) = TX(v) = 0$ and since T is injective, we must have $X(v) = 0$. Thus $T_{t_j-1}^{\gamma,t_j} : V_{t_j-1}^{\gamma,t_j} \to \tilde{V}_{t_j}^{\tilde{\gamma},t_j+1}$ is a linear isomorphism. By applying an appropriate power of $\tilde{X}$, we may conclude that T_i^{γ,t_j} is a linear isomorphism between V_i^{γ,t_j} and $\tilde{V}_{i+1}^{\tilde{\gamma},t_j+1}$ for all $1 \leq j \leq l$ and $-t_j + 1 \leq i \leq t_j - 1$.

We set

$$X_i^{\gamma,t_j} = X|_{V_i^{\gamma,t_j}} \qquad \text{and} \qquad \tilde{X}_i^{\gamma,t_j+1} = \tilde{X}|_{\tilde{V}_i^{\tilde{\gamma},t_j+1}}.$$

Then it is clear that $X_i^{\gamma,t_j} = (T_{-i-2}^{\gamma,t_j})^* T_i^{\gamma,t_j}$ and $\tilde{X}_i^{\tilde{\gamma},t_j+1} = T_{i+1}^{\gamma,t_j}(T_{-i-1}^{\gamma,t_j})^*$. From this we conclude that T is completely determined by the maps $T_{t_j-1}^{\gamma,t_j}$, for $j = 1, \ldots, l$, but now, from equation (5.3), all the $T_{t_j-1}^{\gamma,t_j}$'s are isometries. From this and equation (3.7) it follows immediately that $\mathcal{O}_{\gamma,\tilde{\gamma}}^{\mathrm{Max}}$ is a single $M_X \times \tilde{M}_{\tilde{X}}$-orbit. □

Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \mathfrak{g}'$ be two $\mathfrak{sl}_2$-triples. Given $T \in \mathcal{O}_{\gamma,\tilde{\gamma}}^{\mathrm{Max}}$, we define a map $\phi_T : \tilde{M}_{\tilde{X}} \longrightarrow M_X$ by

$$\phi_T(\tilde{m})v = (T_i^{\gamma,t_j})^{-1} \tilde{m} T_i^{\gamma,t_j} v, \tag{5.6}$$

for all $\tilde{m} \in \tilde{M}_{\tilde{X}}$, and $v \in V_i^{\gamma,t_j}$, where $1 \leq j \leq l$ and $-t_j + 1 \leq i \leq t_j - 1$. Note that as elements of $\mathrm{Hom}(V, \tilde{V})$, we have

$$\tilde{m}T = T\phi_T(\tilde{m}),$$

for all $\tilde{m} \in \tilde{M}_{\tilde{X}}$.

5.4. **Isomorphism of $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ with a symplectic subspace $W_{\gamma,\tilde{\gamma}}$ of $\mathrm{Hom}(V, \tilde{V})$.** Let $\gamma = \{X, H, Y\}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. From the proof of Proposition 5.3, we may assume that

$$V = \bigoplus_{k=-r}^{r} V_k \qquad \text{and} \qquad \tilde{V} = \bigoplus_{k=-r-1}^{r+1} \tilde{V}_k,$$

for some r. We will set

$$W_{\gamma,\tilde{\gamma}} = \bigoplus_{k=-r}^{r} \mathrm{Hom}(V_k, \tilde{V}_k) \subset \mathrm{Hom}(V, \tilde{V}). \tag{5.7}$$

Then it is clear that the restriction of the form $\langle \cdot, \cdot \rangle$, defined in Section 5.1, to $W_{\gamma,\tilde{\gamma}}$ is non-degenerate. Now fix $T \in \mathcal{O}_{\gamma,\tilde{\gamma}}^{\mathrm{Max}}$, and define

$$\begin{array}{rccc} J_T : & \mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1} & \longrightarrow & W_{\gamma,\tilde{\gamma}}, \\ & (R, \tilde{R}) & \mapsto & TR + \tilde{R}T. \end{array} \tag{5.8}$$

Obviously, we have $J_T(R, \tilde{R})(V_k) \subseteq \tilde{V}_k$ for all k.

Lemma 5.8. *J_T defines an isomorphism between $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ and $W_{\gamma,\tilde{\gamma}}$. Furthermore,*

$$\langle J_T(R_1, \tilde{R}_1), J_T(R_2, \tilde{R}_2) \rangle = -\kappa_{-1}(R_1, R_2) + \tilde{\kappa}_{-1}(\tilde{R}_1, \tilde{R}_2),$$

for all R_1, $R_2 \in \mathfrak{g}_{-1}$, $\tilde{R}_1$, $\tilde{R}_2 \in \tilde{\mathfrak{g}}_{-1}$. Here the bilinear form κ on $\mathfrak{g}$ is as in Section 2.2, and likewise for $\tilde{\kappa}$.

Proof. First observe that if R_1, $R_2 \in \mathfrak{g}_{-1}$, then

$$\begin{aligned} \langle TR_1, TR_2 \rangle &= \mathrm{Tr}((TR_1)^* TR_2) = \mathrm{Tr}(R_1^* T^* T R_2) \\ &= -\mathrm{Tr}(R_1 X R_2) = \mathrm{Tr}(R_2 X R_1) \\ &= -\kappa_{-1}(R_1, R_2). \end{aligned}$$

Similarly $\langle \tilde{R}_1 T, \tilde{R}_2 T \rangle = \tilde{\kappa}_{-1}(\tilde{R}_1, \tilde{R}_2)$. On the other hand, if $R \in \mathfrak{g}_{-1}$, $\tilde{R} \in \tilde{\mathfrak{g}}_{-1}$, then

$$\langle TR, \tilde{R}T \rangle = \mathrm{Tr}(R^* T^* \tilde{R} T) = -\mathrm{Tr}(R T^* \tilde{R} T)$$

and

$$\langle \tilde{R}T, TR \rangle = \mathrm{Tr}(T^* \tilde{R}^* T R) = -\mathrm{Tr}(T^* \tilde{R} T R).$$

Therefore, $\langle \tilde{R}T, TR\rangle = 0$, since $\langle\cdot,\cdot\rangle$ is symplectic. We have thus shown that

$$\langle J_T(R_1,\tilde{R}_1), J_T(R_2,\tilde{R}_2)\rangle = -\kappa_{-1}(R_1,R_2) + \tilde{\kappa}_{-1}(\tilde{R}_1,\tilde{R}_2),$$

for all R_1, $R_2 \in \mathfrak{g}_{-1}$, $\tilde{R}_1$, $\tilde{R}_2 \in \tilde{\mathfrak{g}}_{-1}$. But now, since κ_{-1} and $\tilde{\kappa}_{-1}$ are non-degenerate, we conclude that J_T is injective.

Now observe that, for all $S_k \in \operatorname{Hom}(V_k, V_{k-1})$, $S_k - S_k^* \in \mathfrak{g}_{-1}$. On the other hand, we can write $S \in \mathfrak{g}_{-1}$ as the direct sum $\sum_{k=-r}^{r} S_k$, where $S_k \in \operatorname{Hom}(V_k, V_{k-1})$. Then we have that $S_k^* = -S_{-k+1}$ for all k, and hence, the map $S \mapsto S - S^*$ defines a linear isomorphism between $\oplus_{k\le 0}\operatorname{Hom}(V_k, V_{k-1})$ and $\mathfrak{g}_{-1}$. From this, we conclude that

$$\begin{aligned}
\dim \mathfrak{g}_{-1} &= \sum_{k=0}^{-r+1} \dim V_k \cdot \dim V_{k-1} \\
&= \sum_{k=0}^{-r+1} \dim \tilde{V}_{k-1} \cdot \dim V_{k-1} \\
&= \sum_{k=-1}^{-r} \dim \tilde{V}_k \cdot \dim V_k.
\end{aligned}$$

Here we have used the fact that $\dim V_k = \dim \tilde{V}_{k-1}$ for $k \le 0$, in view of the linear isomorphism $T_{-k}^* : \tilde{V}_{k-1} \to V_k$. Similarly,

$$\begin{aligned}
\dim \tilde{\mathfrak{g}}_{-1} &= \sum_{k=0}^{-r} \dim \tilde{V}_k \cdot \dim \tilde{V}_{k-1} \\
&= \sum_{k=0}^{-r} \dim \tilde{V}_k \cdot \dim V_k \\
&= \sum_{k=0}^{r} \dim \tilde{V}_k \cdot \dim V_k.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\dim \mathfrak{g}_{-1} + \dim \tilde{\mathfrak{g}}_{-1} &= \sum_{k=-1}^{-r} \dim \tilde{V}_k \cdot \dim V_k + \sum_{k=0}^{r} \dim \tilde{V}_k \cdot \dim V_k \\
&= \sum_{k=-r}^{r} \dim \operatorname{Hom}(V_k, \tilde{V}_k) = \dim W_{\gamma,\tilde{\gamma}},
\end{aligned}$$

and hence, J_T must be a bijection. □

6. GENERALIZED WHITTAKER MODELS AND HOWE CORRESPONDENCE

6.1. **The smooth oscillator-Heisenberg representation associated to $W_{\gamma,\tilde{\gamma}}$.** Let $H_{\gamma,\tilde{\gamma}}$ be the Heisenberg group associated to the symplectic space $W_{\gamma,\tilde{\gamma}}$. That is, $H_{\gamma,\tilde{\gamma}} = W_{\gamma,\tilde{\gamma}} \times \boldsymbol{k}$, where $\{0\} \times \boldsymbol{k}$ is central, and $(R,0)(S,0) = (R+S, \langle R,S\rangle/2)$. Let $\operatorname{Mp}(W_{\gamma,\tilde{\gamma}})$ be the metaplectic group associated to $W_{\gamma,\tilde{\gamma}}$, and $(\tau_{\gamma,\tilde{\gamma}}, \mathscr{H}_{\gamma,\tilde{\gamma}})$ be the smooth oscillator-Heisenberg representation of $\operatorname{Mp}(W_{\gamma,\tilde{\gamma}}) \ltimes H_{\gamma,\tilde{\gamma}}$ associated to the character ψ.

Recall that we have fixed $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$, and we have an isomorphism J_T from $\mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1}$ to $W_{\gamma,\tilde{\gamma}}$, given in equation (5.8). Define a new invariant bilinear form κ' on $\mathfrak{g}$ by setting $\kappa'(R,S) = -\kappa(R,S)$, for all R, $S \in \mathfrak{g}$. This results in a new symplectic structure on $\mathfrak{g}_{-1}$, given by $\kappa'_{-1}(R,S) = -\kappa_{-1}(R,S)$, for all R, $S \in \mathfrak{g}_{-1}$, and hence a new Heisenberg group H'_{γ}. With this new symplectic structure,

$J_T|_{\mathfrak{g}_{-1}}$ is a morphism of symplectic spaces, and hence, we may extend J_T to an injective morphism of groups $J_T : H'_\gamma \longrightarrow H_{\gamma,\tilde{\gamma}}$. Observe that we can proceed similarly for $\tilde{\mathfrak{g}}_{-1}$, but for this space the modification of the symplectic structure is unnecessary. We obtain in this way a map

$$J_T : H'_\gamma \times H_{\tilde{\gamma}} \longrightarrow H_{\gamma,\tilde{\gamma}}. \tag{6.1}$$

As in Section 3.3, we let $\alpha'_\gamma : N \longrightarrow H'_\gamma$ be the map induced by γ and the bilinear form κ', that is

$$\alpha'_\gamma(\exp R \exp Z) = (R, \kappa'(X, Z)), \quad \text{for all } R \in \mathfrak{g}_{-1},\ Z \in \mathfrak{u}.$$

Similarly, define $\alpha_{\tilde{\gamma}} : \tilde{N} \longrightarrow H_{\tilde{\gamma}}$. By composing with the map J_T, we have a group homomorphism

$$\alpha_T = \alpha_{T,\gamma,\tilde{\gamma}} : N \times \tilde{N} \longrightarrow H_{\gamma,\tilde{\gamma}} \tag{6.2}$$

given by $\alpha_T(n, \tilde{n}) = J_T(\alpha'_\gamma(n), \alpha_{\tilde{\gamma}}(\tilde{n}))$.

Now observe that for all $\tilde{m} \in \tilde{M}_{\tilde{X}}$, $\tilde{R} \in \tilde{\mathfrak{g}}_{-1}$,

$$J_T(\mathrm{Ad}(\tilde{m})\tilde{R}) = \tilde{m}\tilde{R}\tilde{m}^{-1}T = \tilde{m}\tilde{R}T\phi_T(\tilde{m})^{-1}. \tag{6.3}$$

On the other hand, since $\phi_T : \tilde{M}_{\tilde{X}} \longrightarrow M_X$ is surjective, then for any $m \in M_X$ we can find $\tilde{m} \in \tilde{M}_{\tilde{X}}$ such that $\phi_T(\tilde{m}) = m$. Moreover, for all $R \in \mathfrak{g}_{-1}$,

$$J_T(\mathrm{Ad}(\phi_T(\tilde{m}))R) = T\phi(\tilde{m})R\phi(\tilde{m})^{-1} = \tilde{m}TR\phi_T(\tilde{m})^{-1}. \tag{6.4}$$

Using equations (6.3) and (6.4), we define an action of $M_X \times \tilde{M}_{\tilde{X}}$ on $W_{\gamma,\tilde{\gamma}}$, by

$$\tilde{m} \cdot S = \begin{cases} \tilde{m}S\phi_T(\tilde{m})^{-1} & \text{if } S \in J_T(\tilde{\mathfrak{g}}_{-1}) \\ S & \text{if } S \in J_T(\mathfrak{g}_{-1}), \end{cases}$$

if $\tilde{m} \in \tilde{M}_{\tilde{X}}$, and

$$m \cdot S = \begin{cases} S & \text{if } S \in J_T(\tilde{\mathfrak{g}}_{-1}) \\ \tilde{m}S\phi_T(\tilde{m})^{-1} & \text{if } S \in J_T(\mathfrak{g}_{-1}), \end{cases}$$

if $m = \phi_T(\tilde{m}) \in M_X$, for some $\tilde{m} \in \tilde{M}_{\tilde{X}}$. Observe that in particular, for all $\tilde{m} \in \tilde{M}_{\tilde{X}}$, and all $S \in W_{\gamma,\tilde{\gamma}}$, we have

$$(\tilde{m}\phi_T(\tilde{m})) \cdot S = \tilde{m}S\phi_T(\tilde{m})^{-1}. \tag{6.5}$$

Using this action we extend the map α_T to a group homomorphism

$$\alpha_T : M_{\chi_\gamma}N \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N} \longrightarrow \mathrm{Mp}(W_{\gamma,\tilde{\gamma}}) \ltimes H_{\gamma,\tilde{\gamma}}. \tag{6.6}$$

By pulling back, this yields a representation

$$\tau^T_{\gamma,\tilde{\gamma}} := \tau_{\gamma,\tilde{\gamma}} \circ \alpha_T \tag{6.7}$$

of $M_{\chi_\gamma}N \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$ on $\mathscr{H}_{\gamma,\tilde{\gamma}}$. Observe that for all $Z \in \mathfrak{u}$, $v \in \mathscr{H}_{\gamma,\tilde{\gamma}}$,

$$\begin{aligned} \tau^T_{\gamma,\tilde{\gamma}}(\exp Z, \tilde{e})v &= \tau_{\gamma,\tilde{\gamma}}(0, \kappa'(X, Z))v \\ &= \psi(\kappa'(X, Z))v = \psi(-\kappa(X, Z))v \\ &= \chi_{\tilde{\gamma}}(\exp Z)v = \check{\chi}_\gamma(\exp Z)v, \end{aligned}$$

where $\tilde{e}$ is the identity element in $\tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$. Similarly, for all $Z \in \mathfrak{u}$, $v \in \mathscr{H}_{\gamma,\tilde{\gamma}}$,

$$\tau^T_{\gamma,\tilde{\gamma}}(e, \exp \tilde{Z})v = \chi_{\tilde{\gamma}}(\exp \tilde{Z})v,$$

where e is the identity element in $M_{\chi_\gamma}N$. Using this, and the fact that $J_T : \mathfrak{g}_{-1} \oplus \tilde{\mathfrak{g}}_{-1} \to W_{\gamma,\tilde{\gamma}}$ is an isomorphism, we obtain the following result.

Lemma 6.1. *As $M_{\chi_\gamma}N \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$-modules, we have*

$$\tau^T_{\gamma,\tilde{\gamma}} \cong \mathscr{S}_{\check{\chi}_\gamma} \otimes \mathscr{S}_{\chi_{\tilde{\gamma}}},$$

where $\tau^T_{\gamma,\tilde{\gamma}} = \tau_{\gamma,\tilde{\gamma}} \circ \alpha_T$.

6.2. **The main result and the key proposition.** We recall some notation from Section 5. Let (V, B) and $(\tilde{V}, \tilde{B})$ be an ϵ-Hermitian and an $\tilde{\epsilon}$-Hermitian D-module, respectively, with $\varepsilon\tilde{\varepsilon} = -1$. Let $G(V)$, $G(\tilde{V})$ be the corresponding isometry groups and $\mathfrak{g} = \mathfrak{g}(V)$, $\tilde{\mathfrak{g}} = \mathfrak{g}(\tilde{V})$ their Lie algebras. Then $(G(V), G(\tilde{V}))$ form a dual pair in $\mathrm{Sp}(\mathrm{Hom}(V, \tilde{V}))$ in the sense of Howe [7].

Let $\mathrm{Mp}(\mathrm{Hom}(V, \tilde{V}))$ be the metaplectic group. This is the unique topological central extension of $\mathrm{Sp}(\mathrm{Hom}(V, \tilde{V}))$ by $\{\pm 1\}$ such that it splits if $\Bbbk$ is $\mathbb{C}$, and it does not split otherwise. Denote the pullbacks of $G(V)$, $G(\tilde{V})$ in $\mathrm{Mp}(\mathrm{Hom}(V, \tilde{V}))$ by G and $\tilde{G}$, respectively. Then $(G, \tilde{G})$ form a dual pair in $\mathrm{Mp}(\mathrm{Hom}(V, \tilde{V}))$. Let $(\omega, \mathscr{Y})$ be the smooth oscillator representation of $\mathrm{Mp}(\mathrm{Hom}(V, \tilde{V}))$ associated to the character ψ. It is well-known that after tensoring with a genuine character of $G \times \tilde{G}$, $(\omega, \mathscr{Y})$ yields a representation of $G(V) \times G(\tilde{V})$, except when one of them, say V, is an odd dimensional quadratic space and $\Bbbk \neq \mathbb{C}$. In this exceptional case, $\tilde{G}$ is the metaplectic cover of $G(\tilde{V})$, and we shall only consider genuine representations of $\tilde{G}$. For all other cases, representations of $G(V)$ are identified with genuine representations of G (by tensoring with a fixed genuine character of G). With this convention/caveat in mind and for the rest of this article, we will make little distinction between G and $G(V)$, and loosely speak of genuine representations of G. Likewise for $\tilde{G}$ and $G(\tilde{V})$. This convention will lead to significant savings in notation and no confusion is expected for the expert reader.

We state the main result of this article, in a more concrete form than Theorem 1.1 of the introduction.

Theorem 6.2. *Let $(G, \tilde{G})$ be a reductive dual pair, and let $(\omega, \mathscr{Y})$ be the smooth oscillator representation associated to the dual pair $(G, \tilde{G})$ and to the character ψ of $\Bbbk$. Let $(\pi, \mathscr{V})$ be a smooth irreducible genuine representation of G. Let $\mathcal{O} \subset \mathfrak{g}$ be a nilpotent G-orbit in the image of* $\mathrm{Max}\,\mathrm{Hom}(V, \tilde{V})$ *under the moment map φ and let $\tilde{\mathcal{O}} = \Theta(\mathcal{O}) \subset \tilde{\mathfrak{g}}$ be the corresponding nilpotent $\tilde{G}$-orbit. Then*

$$\mathrm{Wh}_{\Theta(\mathcal{O})}(\Theta(\pi)) \cong \mathrm{Wh}_{\mathcal{O}}(\check{\pi}),$$

where $\check{\pi}$ is the contragredient representation of π. More precisely, let γ and $\tilde{\gamma}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. Then, given any $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$, there exists an isomorphism

$$\Phi_T : \mathrm{Wh}_\gamma(\check{\pi}) \longrightarrow \mathrm{Wh}_{\tilde{\gamma}}(\Theta(\pi)),$$

such that

$$\tilde{m}\Phi_T(\lambda) = \Phi_T(\phi_T(\tilde{m})\lambda) \qquad \text{for all } \tilde{m} \in \tilde{M}_{\chi_{\tilde{\gamma}}},\ \lambda \in \mathrm{Wh}_\gamma(\check{\pi}),$$

where $\phi_T : \tilde{M}_{\chi_{\tilde{\gamma}}} \twoheadrightarrow M_{\chi_\gamma}$ is as in equation (5.6).

The key to our proof of Theorem 6.2 is the following result. See Section 4 for the unexplained notation.

Proposition 6.3. *Let $\gamma \subset \mathfrak{g}$ and $\tilde{\gamma} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. Then, given any $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$, there exists a $G \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$-intertwining isomorphism*

$$\Psi_T : \mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}} \longrightarrow \mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}}), \tag{6.8}$$

where $M_{\chi_\gamma}N \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$ acts on $\mathscr{H}_{\gamma,\tilde{\gamma}}$ via the representation $\tau^T_{\gamma,\tilde{\gamma}}$ defined in equation (6.7), and the action of $G \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$ on $\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ is defined in the following way: given $f \in \mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$

and $g \in G$,

$$(g' \cdot f)(g) = f(gg') \qquad \text{for all } g' \in G, \tag{6.9}$$

$$(\tilde{n} \cdot f)(g) = \tau^T_{\gamma,\tilde{\gamma}}(\tilde{n})f(g) \qquad \text{for all } \tilde{n} \in \tilde{N} \tag{6.10}$$

$$(\tilde{m} \cdot f)(g) = \tau^T_{\gamma,\tilde{\gamma}}((\phi_T(\tilde{m}), \tilde{m}))f(\phi_T(\tilde{m})^{-1}g) \qquad \text{for all } \tilde{m} \in \tilde{M}_{\chi_{\tilde{\gamma}}}. \tag{6.11}$$

Before starting with the proof of this proposition, let us show how it implies Theorem 6.2.

Proof (of Theorem 6.2). By Lemma 6.1, we have

$$\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}}) \cong \mathscr{C}(N\backslash G; \mathscr{S}_{\check{\chi}_\gamma} \otimes \mathscr{S}_{\chi_{\tilde{\gamma}}}) \cong \mathscr{C}(N\backslash G; \mathscr{S}_{\check{\chi}_\gamma}) \otimes \mathscr{S}_{\chi_{\tilde{\gamma}}}.$$

Now, given $f \in \mathscr{C}(N\backslash G; \mathscr{S}_{\check{\chi}_\gamma})$ and $v \in \mathscr{S}_{\chi_{\tilde{\gamma}}}$, we identify $f \otimes v$ with an element of $\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ via the above isomorphism. With this identification, we may rewrite equation (6.11) as

$$\begin{aligned} \tilde{m} \cdot (f \otimes v)(g) &= \tau^T_{\gamma,\tilde{\gamma}}(\phi_T(\tilde{m}))f(\phi_T(\tilde{m})^{-1}g) \otimes \tau^T_{\gamma,\tilde{\gamma}}(\tilde{m})v \\ &= (\phi_T(\tilde{m}) \cdot f)(g) \otimes \tau^T_{\gamma,\tilde{\gamma}}(\tilde{m})v, \end{aligned}$$

for all $\tilde{m} \in \tilde{M}_{\chi_{\tilde{\gamma}}}$. See equation (4.11).

By Propositions 6.3 and 4.9, we have the induced isomorphisms:

$$\mathscr{Y}_{(\tilde{U},\chi_{\tilde{\gamma}}),(G,\pi)} \cong \mathscr{C}(N\backslash G; \mathscr{S}_{\check{\chi}_\gamma})_{(G,\pi)} \otimes \mathscr{S}_{\chi_{\tilde{\gamma}}} \cong \pi \otimes W_\gamma(\check{\pi}) \otimes \mathscr{S}_{\chi_{\tilde{\gamma}}},$$

where $\tilde{M}_{\chi_{\tilde{\gamma}}}$ acts on $W_\gamma(\check{\pi})$ through the map $\phi_T : \tilde{M}_{\chi_{\tilde{\gamma}}} \twoheadrightarrow M_{\chi_\gamma}$. On the other hand, we have (from the definition of $\Theta(\pi)$) that

$$\mathscr{Y}_{(G,\pi),(\tilde{U},\chi_{\tilde{\gamma}})} \cong \pi \otimes \Theta(\pi)_{(\tilde{U},\chi_{\tilde{\gamma}})}.$$

Therefore, there exists an isomorphism

$$\Phi_T : \mathrm{Wh}_\gamma(\check{\pi}) \longrightarrow \mathrm{Wh}_{\tilde{\gamma}}(\Theta(\pi)),$$

such that

$$\tilde{m}\Phi_T(\lambda) = \Phi_T(\phi_T(\tilde{m}\lambda)) \qquad \text{for all } \tilde{m} \in \tilde{M}_{\chi_{\tilde{\gamma}}},\ \lambda \in \mathrm{Wh}_\gamma(\check{\pi}).$$

□

In the rest of this section we will closely examine the space $\mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}}$. In fact we will prove a refinement of Proposition 6.3 in which we give an explicit isomorphism Ψ_T in equation (6.8). This is Proposition 6.4. As we have already seen, our main result follows immediately from this proposition.

6.3. **Strategy for the key proposition.** Let $\gamma = \{X, H, Y\} \subset \mathfrak{g}$ and $\tilde{\gamma} = \{\tilde{X}, \tilde{H}, \tilde{Y}\} \subset \tilde{\gamma}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. From the proof of Proposition 5.3, we may assume that

$$V = \bigoplus_{k=-r}^{r} V_k, \qquad \text{and} \qquad \tilde{V} = \bigoplus_{k=-r-1}^{r+1} \tilde{V}_k, \tag{6.12}$$

for some $r \geq 0$. Now fix $T_{\gamma,\tilde{\gamma}} \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$, that is, fix an injective element $T_{\gamma,\tilde{\gamma}} \in \mathrm{Hom}(V, \tilde{V})$, such that $T^*_{\gamma,\tilde{\gamma}}T_{\gamma,\tilde{\gamma}} = X$, $T_{\gamma,\tilde{\gamma}}T^*_{\gamma,\tilde{\gamma}} = \tilde{X}$ and $T_{\gamma,\tilde{\gamma}}(V_k) \subset \tilde{V}_{k+1}$ for all $-r \leq k \leq r$. Then we have a decomposition

$$T_{\gamma,\tilde{\gamma}} = \oplus_{k=-r}^{r} T_k, \tag{6.13}$$

with $T_k \in \mathrm{Hom}(V_k, \tilde{V}_{k+1})$, and T_k is an isomorphism for $k \geq 0$. C.f. Lemma 5.6. Figuratively

$$\begin{tikzcd}[column sep=small] & V_{-r} \arrow[dr, "T_{-r}"] & \oplus V_{-r+1} \arrow[dr, "T_{-r+1}"] & \oplus \cdots \arrow[d, phantom, "\cdots"] & \oplus V_{r-1} \arrow[dr, "T_{r-1}"] & \oplus V_r \arrow[dr, "T_r"] & \\ \tilde{V}_{-r-1} & \oplus \tilde{V}_{-r} & \oplus \tilde{V}_{-r+1} & \oplus \cdots & \oplus \tilde{V}_{r-1} & \oplus \tilde{V}_r & \oplus \tilde{V}_{r+1}. \end{tikzcd}$$

We will use the gradings of V and $\tilde{V}$ associated to H and $\tilde{H}$, respectively, to describe the so-called mixed model of the smooth oscillator representation $(\omega, \mathscr{Y})$ associated to the dual pair $(G, \tilde{G})$. Using this model we will give a convenient description of the space of covariants $\mathscr{Y}_{\tilde{U},\chi_{\tilde{\gamma}}}$ leading to Proposition 6.4. But before we outline the strategy, let us fix some notation. For $l \leq r$ and $m \leq r+1$, set

$$V_{(l)} = \bigoplus_{k=-l}^{l} V_k, \qquad \text{and} \qquad \tilde{V}_{(m)} = \bigoplus_{k=-m}^{m} \tilde{V}_k. \tag{6.14}$$

Let $G_{(l)} = \{g \in G \,|\, g \cdot v = v \text{ for all } v \in V_{(l)}^{\perp}\}$, which is isomorphic with $G(V_{(l)})$. Let $(\omega_{(l),(m)}, \mathscr{Y}_{(l),(m)})$ be the smooth oscillator representation associated to the dual pair $(G_{(l)}, \tilde{G}_{(m)})$ and the character ψ of $\Bbbk$. Set $\mathscr{S}_{(l),m} = \mathscr{S}(\mathrm{Hom}(V_{(l)}, \tilde{V}_m))$, the Schwartz space of $\mathrm{Hom}(V_{(l)}, \tilde{V}_m)$. Similarly, set $\mathscr{S}_{l,(m)} = \mathscr{S}(\mathrm{Hom}(V_l, \tilde{V}_{(m)}))$ and $\mathscr{S}_{l,m} = \mathscr{S}(\mathrm{Hom}(V_l, \tilde{V}_m))$. (Note that $\mathscr{Y}_{(l),(m)}$ can be identified with the Schwartz space of a Lagrangian subspace of $\mathrm{Hom}(V_{(l)}, \tilde{V}_{(m)})$, through the Schrodinger model.) Now observe that

$$\begin{aligned} \mathrm{Hom}(V_{(r)}, \tilde{V}_{(r+1)}) &= \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \oplus \mathrm{Hom}(V_{(r)}, \tilde{V}_{-r-1}) \oplus \mathrm{Hom}(V_{(r)}, \tilde{V}_{(r)}) \\ &= (\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \oplus \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)})) \oplus (\mathrm{Hom}(V_{(r)}, \tilde{V}_{-r-1}) \oplus \mathrm{Hom}(V_r, \tilde{V}_{(r)})) \\ &\quad \oplus \mathrm{Hom}(V_{(r-1)}, \tilde{V}_{(r)}) \end{aligned}$$

and $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \oplus \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)})$, $\mathrm{Hom}(V_{(r)}, \tilde{V}_{-r-1}) \oplus \mathrm{Hom}(V_r, \tilde{V}_{(r)})$ are totally isotropic, complementary subspaces. It then follows, from the standard theory of the oscillator representation [9], that

$$\begin{aligned} \mathscr{Y}_{(r),(r+1)} &\cong \mathscr{S}_{(r),r+1} \otimes \mathscr{Y}_{(r),(r)} \\ &\cong [\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \mathscr{Y}_{(r-1),(r)}, \qquad r > 0. \end{aligned} \tag{6.15}$$

What this equation is saying is that we may interpret the space $\mathscr{Y}_{(r),(r+1)}$ as the space of Schwartz class functions on $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ with values in $\mathscr{Y}_{(r),(r)}$, and the latter, in turn, can be interpreted as the space of Schwartz class functions on $\mathrm{Hom}(V_{-r}, \tilde{V}_{(r)})$ with values in $\mathscr{Y}_{(r-1),(r)}$. More concretely, we describe this space in the following way: given ρ a seminorm on $\mathscr{Y}_{(r),(r)}$, $Z \in \mathscr{D}(\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}))$ ($\mathscr{D}$ denotes the space of constant-coefficient differential operators), $d \in \mathbb{N}$ and $f \in C^\infty(\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}); \mathscr{Y}_{(r),(r)})$, set

$$q_{Z,d,\rho}(f) = \sup_{T \in \mathrm{Hom}(V_{(r)} \tilde{V}_{r+1})} \rho(Zf(T))(1 + \|T\|)^d,$$

where $\|T\|$ is the operator norm of T. Then

$$\mathscr{S}_{(r),r+1} \otimes \mathscr{Y}_{(r),(r)} \cong \left\{ f \in C^\infty(\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}); \mathscr{Y}_{(r),(r)}) \;\middle|\; \begin{array}{c} q_{Z,d,\rho}(f) < \infty, \text{ for all } d \in \mathbb{N}, \\ Z \in \mathscr{D}(\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})), \\ \rho \text{ seminorm on } \mathscr{Y}_{(r),(r)} \end{array} \right\},$$

and likewise for $[\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \mathscr{Y}_{(r-1),(r)}$. Proceeding inductively, we obtain the tensor product decomposition

$$\begin{aligned} \mathscr{Y}_{(r),(r+1)} &\cong [\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \cdots \otimes [\mathscr{S}_{(1),2} \otimes \mathscr{S}_{-1,(1)}] \otimes \mathscr{Y}_{(0),(1)} \\ &\cong [\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \cdots \otimes [\mathscr{S}_{(1),2} \otimes \mathscr{S}_{-1,(1)}] \otimes \mathscr{S}_{(0),1} \otimes \mathscr{Y}_{(0),(0)}. \end{aligned} \tag{6.16}$$

Again we may interpret the space appearing in the right hand side of this equation as the space of Schwartz class functions on $\oplus_{k=r,\ldots,1}[\mathrm{Hom}(V_{(k)}, \tilde{V}_{k+1}) \oplus \mathrm{Hom}(V_{-k}, \tilde{V}_{(k)})] \oplus \mathrm{Hom}(V_{(0)}, \tilde{V}_1)$ with values in $\mathscr{Y}_{(0),(0)}$. This is the mixed model of the smooth oscillator representation associated to the γ,

$\tilde{\gamma}$-gradings. In the rest of this article we will make these identifications of spaces without further explanation.

For simplicity, assume for the moment that $\mathcal{O}$ and $\Theta(\mathcal{O})$ are even orbits, that is $\mathfrak{g}_{-1} = 0$ and $\tilde{\mathfrak{g}}_{-1} = 0$. (By Section 5.4, this is equivalent to $W_{\gamma,\tilde{\gamma}} = \bigoplus_{k=-r}^{r} \mathrm{Hom}(V_k, \tilde{V}_k) = 0$.) In this case we have that $\mathscr{Y}_{(0),(0)} = \mathbb{C}$ and we may define a continuous linear functional $\lambda_{(r)} \in \mathscr{Y}'_{(r),(r+1)}$ by setting

$$\lambda_{(r)}(f) = f(T_r, T_{-r}, \ldots, T_1, T_{-1}, T_0),$$

where $f \in \mathscr{Y}_{(r),(r+1)} \cong [\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \cdots \otimes [\mathscr{S}_{(1),2} \otimes \mathscr{S}_{-1,(1)}] \otimes \mathscr{S}_{(0),1} \otimes \mathscr{Y}_{(0),(0)}$. As we will see later in this section, for such an f we have that

$$\lambda_{(r)}(\omega_{(r),(r+1)}(n)f) = \chi_{\gamma}^{-1}(n)\lambda_{(r)}(f) \qquad \text{for all } n \in U = N,$$

and

$$\lambda_{(r)}(\omega_{(r),(r+1)}(\tilde{n})f) = \chi_{\tilde{\gamma}}(\tilde{n})\lambda_{(r)}(f) \qquad \text{for all } \tilde{n} \in \tilde{U} = \tilde{N}.$$

Now given $f \in \mathscr{Y}_{(r),(r+1)}$, set

$$f_{(r)}(g) = \lambda_{(r)}(\omega_{(r),(r+1)}(g)f),$$

for $g \in G$. It is then immediate that $f_{(r)} \in C^\infty(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ but we will show that actually $f_{(r)} \in \mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$. (Here we note that $\mathscr{H}_{\gamma,\tilde{\gamma}}$ is 1-dimensional since $W_{\gamma,\tilde{\gamma}} = 0$.) On the other hand, since M_χ and $\tilde{M}_{\chi_{\tilde{\gamma}}}$ preserve the gradings on V and $\tilde{V}$, respectively, it is straightforward to check in this case that

$$(\omega_{(r),(r+1)}(\tilde{m})f)_{(r)}(g) = f_{(r)}(\phi_T(\tilde{m})^{-1}g), \text{for all } f \in \mathscr{Y}_{(r),(r+1)},\ \tilde{m} \in \tilde{M}_{\chi_{\tilde{\gamma}}},\ g \in G.$$

It then follows that the map $f \mapsto f_{(r)}$ induces a $G \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$-intertwining map

$$\Psi_T : (\mathscr{Y}_{(r),(r+1)})_{\tilde{U},\chi_{\tilde{\gamma}}} \longrightarrow \mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}}),$$

that we will show to be an isomorphism.

To describe the map Ψ_T in the general case, observe that if $(\tau_{\gamma,\tilde{\gamma}}, \mathscr{H}_{\gamma,\tilde{\gamma}})$ is the smooth Heisenberg representation associated to $W_{\gamma,\tilde{\gamma}}$ (as in Section 6.1), then, following arguments similar to those leading to equation (6.16), we obtain the following tensor product decomposition:

$$\mathscr{H}_{\gamma,\tilde{\gamma}} \cong \mathscr{S}_{-r,-r} \otimes \mathscr{S}_{-r+1,-r+1} \cdots \otimes \mathscr{S}_{-1,-1} \otimes \mathscr{Y}_{(0),(0)}. \tag{6.17}$$

Again we may identify the space $\mathscr{H}_{\gamma,\tilde{\gamma}}$ with the space of Schwartz class functions on $\mathrm{Hom}(V_{-r}, \tilde{V}_{-r}) \oplus \cdots \oplus \mathrm{Hom}(V_{-1}, \tilde{V}_{-1})$ with values in $\mathscr{Y}_{(0),(0)}$. Now, given $f \in \mathscr{Y}_{(r),(r+1)}$, set

$$f_{(r)}(g)(S_{-r}, \ldots, S_{-1}) = (\omega_{(r),(r+1)}(g)f)(T_r, T_{-r} + S_{-r}, \ldots, T_1, T_{-1} + S_{-1}, T_0), \tag{6.18}$$

for all $g \in G$, $S_{-k} \in \mathrm{Hom}(V_{-k}, \tilde{V}_{-k})$, $k = 1, \ldots, r$. Note that when $\mathcal{O}$ and $\Theta(\mathcal{O})$ are both even this definition agrees with the one given before.

The following result is a refinement of Proposition 6.3.

Proposition 6.4. *Let $\gamma \subset \mathfrak{g}$ and $\tilde{\gamma} \subset \tilde{\mathfrak{g}}$ be two $\mathfrak{sl}_2$-triples of type $\mathcal{O}$ and $\Theta(\mathcal{O})$, respectively. Fix $T \in \mathcal{O}^{\mathrm{Max}}_{\gamma,\tilde{\gamma}}$. Given $f \in \mathscr{Y}_{(r),(r+1)}$, define $f_{(r)} \in C^\infty(G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ as in equation (6.18). Then the map $f \mapsto f_{(r)}$ induces a $G \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$-intertwining isomorphism*

$$\Psi_T : (\mathscr{Y}_{(r),(r+1)})_{\tilde{U},\chi_{\tilde{\gamma}}} \longrightarrow \mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}}).$$

where $G \times \tilde{M}_{\chi_{\tilde{\gamma}}}\tilde{N}$ acts on $\mathscr{C}(N\backslash G; \mathscr{H}_{\gamma,\tilde{\gamma}})$ via equations (6.9)–(6.11).

As we have already shown, the existence of such a Ψ_T is enough to prove Theorem 6.2. Before proceeding to the proof of this key proposition, we introduce all the remaining notation.

Recall that we have set $G_{(l)} = \{g \in G \,|\, g \cdot v = v \text{ for all } v \in V_{(l)}^{\perp}\}$. Let P_l be the stabilizer of V_{-l} in $G_{(l)}$. Then $P_l = M_{(l)}N_l$, where N_l is the unipotent radical of P_l, $M_{(l)} = M_l \times G_{(l-1)}$, and

$M_l \cong \mathrm{GL}(V_{-l})$. Now observe that, if $S \in \mathrm{Hom}(V_{(l-1)}, V_{-l})$, then $S - S^* \in \mathfrak{n}_l$. Hence, we can use the map $S \mapsto S - S^*$ to define a Lie algebra isomorphism

$$\mathrm{Hom}(V_{(l-1)}, V_{-l}) \oplus \mathfrak{z}_l \cong \mathfrak{n}_l, \tag{6.19}$$

where $\mathfrak{z}_l = \{Z \in \mathrm{Hom}(V_l, V_{-l}) \,|\, Z^* = -Z\}$ is central, and $[T, S] = ST^* - TS^*$ for all T, $S \in \mathrm{Hom}(V_{(l-1)}, V_{-l})$. In what follows we will make implicit use of this isomorphism to identify these two spaces.

In a similar fashion, we define $\tilde{G}_{(m)} \cong G(\tilde{V}_{(m)})$, and observe that if we set $\tilde{P}_m$ to be the stabilizer of $\tilde{V}_m$ in $\tilde{G}_{(m)}$, then $\tilde{P}_m = \tilde{M}_{(m)}\tilde{N}_m$, where $\tilde{N}_m$ is the unipotent radical of $\tilde{P}_m$, $\tilde{M}_{(m)} = \tilde{M}_m \times \tilde{G}_{(m-1)}$, and $\tilde{M}_m \cong \mathrm{GL}(\tilde{V}_m)$. Similarly to (6.19), we use the map $\tilde{S} \mapsto \tilde{S} - \tilde{S}^*$ from $\mathrm{Hom}(\tilde{V}_m, \tilde{V}_{(m-1)})$ to $\tilde{\mathfrak{n}}_m$ to define a Lie algebra isomorphism

$$\mathrm{Hom}(\tilde{V}_m, \tilde{V}_{(m-1)}) \oplus \tilde{\mathfrak{z}}_m \cong \tilde{\mathfrak{n}}_m, \tag{6.20}$$

where, again, $\tilde{\mathfrak{z}}_m = \{\tilde{Z} \in \mathrm{Hom}(\tilde{V}_m, \tilde{V}_{-m}) \,|\, \tilde{Z}^* = -\tilde{Z}\}$ is central, and $[\tilde{T}, \tilde{S}] = \tilde{S}^*\tilde{T} - \tilde{T}^*\tilde{S}$ for all $\tilde{T}$, $\tilde{S} \in \mathrm{Hom}(\tilde{V}_m, \tilde{V}_{(m-1)})$. We will also set

$$N_{(l)} = N \cap G_{(l)}, \qquad U_l = U \cap N_l, \qquad \text{and} \qquad U_{(l)} = U \cap N_{(l)}, \tag{6.21}$$

with similar definitions for $\tilde{N}_{(m)}$, $\tilde{U}_m$ and $\tilde{U}_{(m)}$. Let $\chi = \chi_\gamma$ be the character of U associated to the $\mathfrak{sl}_2$-triple γ, and let $\chi_l = \chi|_{U_l}$, $\chi_{(l)} = \chi|_{U_{(l)}}$. Similarly we define $\tilde{\chi} = \chi_{\tilde{\gamma}}$, $\tilde{\chi}_m$ and $\tilde{\chi}_{(m)}$. Finally, define $X_{(l)} \in \mathrm{End}(V_{(l)})$ in the following way: for $k < l-1$ we define $X_{(l)}|_{V_k} = X|_{V_k}$ and for $l - 1 \le k \le l$ set $X_{(l)}|_{V_k} = 0$. Then it is clear that $X_{(l)} \in \mathfrak{g}_{(l)} := \mathfrak{g}(V_{(l)})$ is a nilpotent element. Let $H_{(l)} = H|_{V_{(l)}}$. Then $H_{(l)} \in \mathfrak{g}_{(l)}$ is semisimple, and $[H_{(l)}, X_{(l)}] = 2X_{(l)}$. Therefore, there exists $Y_{(l)} \in \mathfrak{g}_{(l)}$ such that $\gamma_{(l)} = \{X_{(l)}, H_{(l)}, Y_{(l)}\}$ is an $\mathfrak{sl}_2$-triple. Observe that the parabolic subgroup associated to this $\mathfrak{sl}_2$-triple is precisely $P_{\gamma_{(l)}} = P_\gamma \cap G_{(l)} = M_{(l)}N_{(l)}$, where $M_{(l)} = M \cap G_{(l)}$. Furthermore, observe that $\chi_{\gamma_{(l)}} = \chi_{(l)}$, and hence $M_{\chi_{\gamma_{(l)}}} = M_{\chi_{(l)}}$. We make analogous definitions for $\tilde{X}_{(l)}$, $\tilde{\gamma}_{(l)}$, etc.

We recall the following explicit formulas for $\omega_{(r),(r+1)}|_{G \times \tilde{P}_{r+1}}$ [9]: for $f \in \mathscr{Y}_{(r),(r+1)} \cong \mathscr{S}_{(r),r+1} \otimes \mathscr{Y}_{(r),(r)}$ and $T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$,

$$(\omega_{(r),(r+1)}(g)f)(T) = \omega_{(r),(r)}(g)[f(Tg)] \qquad \text{for all } g \in G = G(V_{(r)}), \tag{6.22}$$

$$(\omega_{(r),(r+1)}(\tilde{m})f)(T) = \nu(\tilde{m})[f(\tilde{m}^{-1}T)] \qquad \text{for all } \tilde{m} \in \tilde{M}_{r+1} \cong \mathrm{GL}(\tilde{V}_{r+1}), \tag{6.23}$$

$$(\omega_{(r),(r+1)}(\tilde{g})f)(T) = \omega_{(r),(r)}(\tilde{g})[f(T)] \qquad \text{for all } \tilde{g} \in \tilde{G}_{(r)} \cong G(\tilde{V}_{(r)}), \tag{6.24}$$

$$(\omega_{(r),(r+1)}(\exp \tilde{Z})f)(T) = \psi(\mathrm{Tr}\, \tilde{Z}TT^*/2)[f(T)] \qquad \text{for all } \tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}, \tag{6.25}$$

$$(\omega_{(r),(r+1)}(\exp \tilde{R})f)(T) = \omega_{(r),(r)}(-\tilde{R}T)[f(T)] \qquad \text{for all } \tilde{R} \in \mathrm{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)}). \tag{6.26}$$

Here ν is a certain character whose explicit form will not concern us. In the last equation we implicitly understand that $\mathscr{Y}_{(r),(r)}$ carries an action of the Heisenberg group associated to $\mathrm{Hom}(V_{(r)}, \tilde{V}_{(r)})$. We also observe the following: since $\tilde{V}_{r+1} \subset \tilde{V}_{(r+1)}$ is stable under the action of $\tilde{P}_{r+1}$, we may consider $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ as a $G \times \tilde{P}_{r+1}$-module with trivial $\tilde{G}_{(r)} \times \tilde{N}_{r+1}$ ($\subset \tilde{P}_{r+1}$) action. From equations (6.22)–(6.26), it is then clear that the actions $\omega_{(r),(r+1)}|_{G \times \tilde{P}_{r+1}}$ may be interpreted as coming from the usual vector bundle action of $G \times \tilde{P}_{r+1}$ on the trivial vector bundle with base $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ and the fiber $\mathscr{Y}_{(r),(r)}$. (This action depends on T.) To be more precise, we may rewrite equations (6.22)–(6.26) in terms of an action σ_T of $G \times \tilde{P}_{r+1}$ on $\mathscr{Y}_{(r),(r)}$, as follows:

$$(\omega_{(r),(r+1)}(g)f)(T) = \sigma_T(g)f(Tg) \qquad \text{for all } g \in G \tag{6.27}$$

$$(\omega_{(r),(r+1)}(\tilde{p})f)(T) = \sigma_T(\tilde{p})f(\tilde{p}^{-1}T) \qquad \text{for all } \tilde{p} \in \tilde{P}_{r+1}, \tag{6.28}$$

where $\sigma_T(g) = \omega_{(r),(r)}(g)$ for all $g \in G$, and

$$\sigma_T(\tilde{p}) = \psi(\operatorname{Tr} \tilde{Z} T T^*/2)\nu(\tilde{m})\omega_{(r),(r)}(\tilde{g})\omega_{(r),(r)}(\tilde{R} T),$$

for $\tilde{p} = \tilde{m}\tilde{g}(\exp \tilde{R})(\exp \tilde{Z})$, with $\tilde{m} \in \tilde{M}_{r+1}$, $\tilde{g} \in \tilde{G}_{(r)}$, $\tilde{R} \in \operatorname{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)})$, $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$.

6.4. **The inductive step: relating covariants of $\mathscr{Y}_{(r),(r+1)}$ with $\mathscr{Y}_{(r-1),(r)}$.** Given a function $f \in \mathscr{Y}_{(r),(r+1)} \cong [\mathscr{S}_{(r),r+1} \otimes \mathscr{S}_{-r,(r)}] \otimes \mathscr{Y}_{(r-1),(r)}$, define a new function $f_r \in C^\infty(G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$ by

$$f_r(g)(S) = [\omega_{(r),(r+1)}(g)f](T_r, T_{-r} + S), \qquad \text{for all } g \in G,\ S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{-r}), \tag{6.29}$$

where $T_{\gamma,\tilde{\gamma}} = \sum_{k=-r}^{r} T_k$ is as before. Then, from equations (6.25) and (6.26), we have that for all $f \in \mathscr{Y}_{(r),(r+1)}$ $g \in G$, $S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{-r})$,

$$\begin{aligned}
(\omega_{(r),(r+1)}(\exp \tilde{Z})f)_r(g)(S) &= (\omega_{(r),(r+1)}(\exp \tilde{Z})[\omega_{(r),(r+1)}(g)f])(T_r, T_{-r} + S) \\
&= \psi(\operatorname{Tr} \tilde{Z} T_r T_r^*/2)(\omega_{(r),(r+1)}(g)f)(T_r, T_{-r} + S) \\
&= f_r(g)(S), \qquad\qquad\qquad\qquad\qquad\qquad \text{(6.30)}
\end{aligned}$$

for all $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$, and

$$\begin{aligned}
(\omega_{(r),(r+1)}(\exp \tilde{R})f)_r(g)(S) &= (\omega_{(r),(r+1)}(\exp \tilde{R})[\omega_{(r),(r+1)}(g)f])(T_r, T_{-r} + S) \\
&= (\omega_{(r),(r)}(-\tilde{R} T_r)[\omega_{(r),(r+1)}(g)f](T_r))(T_{-r} + S) \\
&= \psi(\operatorname{Tr}((T_{-r} + S)^* \tilde{R} T_r))[\omega_{(r),(r+1)}(g)f](T_r, T_{-r} + S) \\
&= \psi(\operatorname{Tr}(T_{-r}^* \tilde{R} T_r))[\omega_{(r),(r+1)}(g)f](T_r, T_{-r} + S) \\
&= \chi_{\tilde{\gamma}}(\exp \tilde{R}) f_r(g)(S), \qquad\qquad\qquad\qquad \text{(6.31)}
\end{aligned}$$

for all $\tilde{R} \in \operatorname{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r-1)} \oplus \tilde{V}_{-r}) \subset \tilde{\mathfrak{u}}_{r+1}$, in other words, for all $\tilde{u} \in \tilde{U}_{r+1}$,

$$(\omega_{(r),(r+1)}(\tilde{u})f)_r(g)(S) = \chi_{\tilde{\gamma}}(\tilde{u}) f_r(g)(S).$$

It is then clear that understanding the map $f \mapsto f_r$ is an important first step towards understanding the more complicated map $f \mapsto f_{(r)}$.

We start by observing that $G_{(r-1)}N_r$ is the stabilizer of T_r in G. Hence, according to equation (6.27) we obtain a representation $(\sigma_{T_r}, \mathscr{Y}_{(r),(r)})$ of $G_{(r-1)}N_r$, where we are viewing $\mathscr{Y}_{(r),(r)}$ as the fiber over the point $T_r \in \operatorname{Hom}(V_{(r)}, \tilde{V}_{r+1})$. Now, since by definition $\sigma_{T_r}(g)\varphi = \omega_{(r),(r)}(g)\varphi$ for all $\varphi \in \mathscr{Y}_{(r),(r)}$, we may use the formulas given in [9] to describe this action explicitly: given $\varphi \in \mathscr{Y}_{(r),(r)}$, $S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{(r)})$,

$$\begin{aligned}
(\sigma_{T_r}(\exp Z)\varphi)(S) &= \psi(\operatorname{Tr} S^* S Z/2)\varphi(S), && \text{for all } Z \in \mathfrak{z}_r, && \text{(6.32)} \\
(\sigma_{T_r}(\exp R)\varphi)(S) &= \omega_{(r-1),(r)}(SR)[\varphi(S)], && \text{for all } R \in \operatorname{Hom}(V_{(r-1)}, V_{-r}), && \text{(6.33)} \\
(\sigma_{T_r}(g)\varphi)(S) &= \omega_{(r-1),(r)}(g)[\varphi(S)], && \text{for all } g \in G_{(r-1)}. && \text{(6.34)}
\end{aligned}$$

Now, given $\varphi \in \mathscr{Y}_{(r),(r)}$, set $\bar{\varphi}(S) = \varphi(T_{-r} + S)$ for all $S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{-r})$. It is clear that the map $\varphi \mapsto \bar{\varphi}$ defines a surjection

$$^{-}\colon\ \mathscr{Y}_{(r),(r)} \twoheadrightarrow \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}.$$

Furthermore, using equations (6.32)–(6.34), we may thus define a representation $(\bar{\sigma}_{T_r}, \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$ of $G_{(r-1)}N_r$ explicitly by the following formulas: given $\varphi \in \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}$, $S \in$

$\mathrm{Hom}(V_{-r}, \tilde{V}_{-r})$,

$$\begin{aligned}
&(6.35) \quad (\bar{\sigma}_{T_r}(\exp Z)\varphi)(S) &&= \psi(\mathrm{Tr}(T_{-r}+S)^*(T_{-r}+S)Z/2)\varphi(S) = \varphi(S), \qquad \text{for } Z \in \mathfrak{z}_r,\\
&(6.36) \quad (\bar{\sigma}_{T_r}(\exp R)\varphi)(S) &&= \omega_{(r-1),(r)}((T_{-r}+S)R)[\varphi(S)], \qquad \text{for } R \in \mathrm{Hom}(V_{(r-1)}, V_{-r}),\\
&(6.37) \qquad\quad (\bar{\sigma}_{T_r}(g)\varphi)(S) &&= \omega_{(r-1),(r)}(g)[\varphi(S)], \qquad \text{for } g \in G_{(r-1)}.
\end{aligned}$$

Observe that if $f \in \mathscr{Y}_{(r),(r+1)}$, then $f_r(g) = \overline{[\omega_{(r),(r+1)}(g)f](T_r)}$. It follows immediately that

$$f_r \in C^\infty(G_{(r-1)}N_r\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}).$$

Now, given ρ a seminorm in $\mathscr{Y}_{(r-1),(r)}$, $Z_1 \in \mathscr{D}(\mathrm{Hom}(V_{-r}, \tilde{V}_{-r}))$, $Z_2 \in U(\mathfrak{g})$, d_1, $d_2 \in \mathbb{N}$ and $f \in C^\infty(G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$, let

$$q_{Z_1,Z_2,d_1,d_2,\rho}(f) = \sup_{\substack{k \in K,\, m \in M_r\\ S \in \mathrm{Hom}(V_{-r}, \tilde{V}_{-r})}} \rho(Z_1[Z_2 f(mk)](S))\|m\|^{d_1}(1+\|S\|)^{d_2}, \tag{6.38}$$

where $K \subset G$ is a maximal compact subgroup as in Section 4.1. Set

$$\begin{aligned}
&\mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}) =\\
&\qquad \left\{ f \in C^\infty(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}) \;\middle|\; \begin{array}{c} q_{Z_1,Z_2,d_1,d_2,\rho}(f) < \infty, \text{ for all}\\ q_{Z_1,Z_2,d_1,d_2,\rho} \text{ as in (6.38)} \end{array} \right\}.
\end{aligned}$$

Lemma 6.5. *The map $f \mapsto f_r$ induces a G-intertwining isomorphism*

$$\Psi_r : (\mathscr{Y}_{(r),(r+1)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} \longrightarrow \mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}),$$

where $N_r G_{(r-1)}$ acts on $\mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}$ by the representation $\bar{\sigma}_{T_r}$, given in equations (6.35)–(6.37).

Proof. Let $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$. As in the Appendix, we will identify λ with the $\mathscr{Y}'_{(r),(r)}$-valued distribution on $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ that sends $f \in \mathscr{S}_{(r),r+1}$ to the linear functional $v \mapsto \lambda(f \otimes v)$. In other words, and with some abuse of notation, we set

$$\lambda(f)(v) := \lambda(f \otimes v) \qquad \text{for all } f \in \mathscr{S}_{(r),r+1},\ v \in \mathscr{Y}_{(r),(r)}.$$

Now recall that for $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$, $f \in \mathscr{S}_{(r),r+1}$ and $T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$, we have $\omega_{(r),(r+1)}(\exp \tilde{Z})f(T) = \psi(\mathrm{Tr}\, \tilde{Z}TT^*/2)f(T)$. Hence, for all such $\tilde{Z}$ we have that

$$\omega_{(r),(r+1)}(\exp \tilde{Z})\lambda = \psi(\mathrm{Tr}\, \tilde{Z}TT^*/2)\lambda, \tag{6.39}$$

that is, the action of $\exp \tilde{Z}$ on λ is given by multiplication by the function $T \mapsto \psi(\mathrm{Tr}\, \tilde{Z}TT^*/2)$. On the other hand, since $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ and since $\tilde{\mathfrak{z}}_{r+1} \subseteq \mathfrak{u}_{r+1}$, we must have that

$$\omega_{(r),(r+1)}(\exp \tilde{Z})\lambda = \psi(\mathrm{Tr}\, \tilde{Z}\tilde{X}/2)\lambda = \lambda, \tag{6.40}$$

where the latter equality is due to the fact that $\tilde{Z} \in \mathfrak{g}_{-2r-2}(\tilde{V})$, $\tilde{X} \in \mathfrak{g}_2(\tilde{V})$ and $r > 0$. Define $h_{\tilde{Z}} \in C^\infty(\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}))$ by $h_{\tilde{Z}}(T) = \psi(\mathrm{Tr}\, \tilde{Z}TT^*/2) - 1$ for all $T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$. Then we can reformulate equations (6.39) and (6.40) by saying that $h_{\tilde{Z}}\lambda = 0$, for all $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$. It then follows immediately that

$$\begin{aligned}
\mathrm{supp}\, \lambda &\subseteq \bigcap_{\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}} \{T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \,|\, h_{\tilde{Z}}(T) = 0\}\\
&= \{T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \,|\, TT^* = 0\}.
\end{aligned}$$

Let

$$\mathrm{Hom}_{NM}(V_{(r)}, \widetilde{V}_{r+1}) = \{T \in \mathrm{Hom}(V_{(r)}, \widetilde{V}_{r+1}) \,|\, TT^* = 0\}, \quad \text{and} \tag{6.41}$$

$$\mathrm{Hom}_{GNM}(V_{(r)}, \widetilde{V}_{r+1}) = \{T \in \mathrm{Hom}(V_{(r)}, \widetilde{V}_{r+1}) \mid TT^* = 0 \text{ and } T \text{ has maximal rank}\}. \tag{6.42}$$

(Here the subscripts MN and GMN stand for null mappings and generic null mappings, respectively.)

We claim that the natural map

$$\mathscr{Y}_{(r),(r+1)} \twoheadrightarrow (\mathscr{Y}_{(r),(r+1)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$$

factors through the intermediate space $\mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)}$. In other words, if $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, then we can identify λ with a $\mathscr{Y}'_{(r),(r)}$-valued tempered distribution *living* on $\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})$. So far we have only shown that if $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, then $\operatorname{supp}\lambda \subset \mathrm{Hom}_{NM}(V_{(r)}, \tilde{V}_{r+1})$. Let

$$\mathcal{A} = \{T \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1}) \mid T \text{ is singular}\}.$$

Clearly $\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1}) = \mathrm{Hom}_{NM}(V_{(r)}, \tilde{V}_{r+1}) \cap \mathcal{A}^c$, where $\mathcal{A}^c$ is the complement of $\mathcal{A}$. So, as a first step towards proving our claim, we will show that

$$\text{if } \lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} \quad \text{and} \quad \operatorname{supp}\lambda \subset \mathcal{A}, \qquad \text{then} \quad \lambda = 0. \tag{6.43}$$

Observe that $\mathcal{A} \subset \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ is invariant under the natural action of $G(V_{(r)}) \times \mathrm{GL}(\tilde{V}_{r+1})$ and decomposes as a finite union of orbits. (Actually, $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ itself decomposes as a finite union of orbits.) Furthermore, if $\mathcal{C} \subset \mathcal{A}$ is a closed invariant subset, then there exists an orbit $\mathcal{O} \subset \mathcal{C}$ that is open in $\mathcal{C}$. Therefore, we may use a Bruhat type argument to prove statement (6.43). To be precise, we will show that if $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, $\mathcal{C} \subset \mathcal{A}$ is a non-empty closed invariant subset containing $\operatorname{supp}\lambda$ and $\mathcal{O} \subset \mathcal{C}$ is a relatively open orbit, then λ vanishes in the complement of $\mathcal{C}\backslash\mathcal{O}$. Once this is done, a straightforward inductive argument finishes the proof of statement (6.43).

If $\mathfrak{k}$ is non-Archimedian then we may immediately identify λ with a distribution living on $\mathcal{C}$. In this case, by restricting λ to $\mathcal{O}$ we obtain a well-defined distribution on this orbit. So we will just need to focus on the case where $\mathfrak{k}$ is Archimedian. Given any $x \in \mathcal{O}$, there exists a neighborhood $U_x \subset \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ of x such that the order of $\lambda|_{U_x}$ is $\leq l$ for some $l \in \mathbb{N}$. Therefore, we may find a locally finite open cover $\{U_i\}_{i\in\mathbb{N}}$ of $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ and a partition of unity $\{\rho_i\}$ subordinated to $\{U_i\}$ such that $\lambda = \sum_i \rho_i\lambda$ and each $\rho_i\lambda$ is of order $\leq l_i$, for some $l_i \in \mathbb{N}$. Now, since multiplication by a C^∞-function clearly commutes with the action of $\tilde{U}_{r+1}$, we have that, if $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, then $\rho_i\lambda$ is also in $(\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ for all i. The upshot is that in order to prove statement (6.43) it is enough to just consider distributions $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ of finite order. But in such a case, to prove the vanishing of λ in the complement of $\mathcal{C}\backslash\mathcal{O}$, it suffices to show that $D'(\mathcal{O}; \mathscr{Y}_{(r),(r)} \otimes M^{(l)'})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} = 0$ for all $l \geq 0$ [14]. Here $M^{(l)}$ is the l-th transverse jet bundle of $\mathcal{O}$ (see the Appendix). Now observe that, given $S \in \mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$ and $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$,

$$\begin{aligned}
\frac{\partial}{\partial S}(\exp \tilde{Z}) \cdot f(T) &= \left.\frac{d}{dt}\right|_{t=0} \psi(\mathrm{Tr}(T+tS)(T+tS)^*\tilde{Z}/2) f(T+tS) \\
&= \psi(\mathrm{Tr}\, TT^*\tilde{Z}/2)\psi(\mathrm{Tr}\, ST^*\tilde{Z}/2) f(T) + \psi(\mathrm{Tr}\, TT^*\tilde{Z}/2)\psi(\mathrm{Tr}\, TS^*\tilde{Z}/2) f(T) \\
&\quad + \psi(\mathrm{Tr}\, TT^*\tilde{Z}/2)\frac{\partial f}{\partial S}(T),
\end{aligned}$$

and if $R \in \mathrm{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)}) \subset \tilde{\mathfrak{n}}_{r+1}$, then

$$\begin{aligned}\frac{\partial}{\partial S}(\exp R)\cdot f(T) &= \left.\frac{d}{dt}\right|_{t=0} \omega_{(r),(r)}(R(T+tS))f(T+tS)\\ &= \left.\frac{d}{dt}\right|_{t=0} \omega_{(r),(r)}(RT)\omega_{(r),(r)}(RtS)f(T+tS)\\ &= \omega_{(r),(r)}(RT)d\omega_{(r),(r)}(RS)f(T)+\omega_{(r),(r)}(RT)\frac{\partial}{\partial S}f(T).\end{aligned}$$

What the above couple of equations imply is that the action of $\tilde{N}_{r+1}$ (and in particular the action of $\tilde{U}_{r+1}$) on the l-th transverse bundle is trivial, and hence $D'(\mathcal{O};\mathscr{Y}_{(r),(r)}\otimes (M^{(l)})')^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} \cong M^{(l)}\otimes D'(\mathcal{O};\mathscr{Y}_{(r),(r)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$. From all this we see that, in order to prove statement (6.43), it suffices to show that

$$D'(\mathcal{O};\mathscr{Y}_{(r),(r)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} = 0 \tag{6.44}$$

for every orbit $\mathcal{O}\subset\mathcal{A}$.

At this point we come back to the general case, where $\mathfrak{k}$ can be Archimedean or non-Archimedean. Let $\lambda \in D'(\mathcal{O};\mathscr{Y}_{(r),(r)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, where $\mathcal{O}\subset\mathcal{A}$ is a $G(V_{(r)})\times \mathrm{GL}(\tilde{V}_{r+1})$-orbit. As we have already seen, the equivariance of λ implies that $\mathrm{supp}\,\lambda \in \mathcal{O}\cap \mathrm{Hom}_{NM}(V_{(r)},\tilde{V}_{r+1})$. Since the latter set is $G(V_{(r)})\times\mathrm{GL}(\tilde{V}_{r+1})$ invariant, we see that actually $\mathcal{O}\subset \mathrm{Hom}_{NM}(V_{(r)},\tilde{V}_{r+1})$. From equations (6.27) and (6.28), it is clear that once we fix any $T\in\mathcal{O}$ we could describe $C_c^\infty(\mathcal{O})\otimes\mathscr{Y}_{(r),(r)}$ as an induced representation. (Here we are using the identifications $G\cong G(V_{(r)})$ and $\tilde{M}_{r+1}\cong \mathrm{GL}(\tilde{V}_{r+1})$.) Translating under the action of $G(V_{(r)})$, if necessary, we may assume that $T\in\mathcal{O}$ satisfies $T|_{\oplus_{k=-r}^{r-1}V_k} = 0$. To see why, let $T\in \mathrm{Hom}_{NM}(V_{(r)},\tilde{V}_{r+1})$ $(\subseteq \mathrm{Hom}_{NM}(V_{(r)},\tilde{V}_{(r+1)}))$, and so $T^*|_{\tilde{V}_{r+1}^{\perp}} = 0$. Since $\tilde{V}_{r+1}^{\perp} = \tilde{V}_{(r)}\oplus\tilde{V}_{r+1}$ we have that $\mathrm{Im}\,T^*$ $(\subset V_{(r)})$ is a totally isotropic subspace with dimension less than or equal to $\dim\tilde{V}_{-r-1} = \dim V_{-r}$. We may thus assume, by translating under $G(V_{(r)})$ if necessary, that $\mathrm{Im}\,T^*\subset V_{-r}$, which will imply that $T|_{\oplus_{k=-r}^{r-1}V_k} = 0$. We identify such a T with an element of $\mathrm{Hom}(V_r,\tilde{V}_{r+1})\subset \mathrm{Hom}(V_{(r)},\tilde{V}_{r+1})$ in an obvious way. Now we have an isomorphism of $G\times\tilde{P}_{r+1}$-modules:

$$C_c^\infty(\mathcal{O})\otimes\mathscr{Y}_{(r),(r)} \cong C_c^\infty - \mathrm{Ind}_{\mathrm{Stab}_T}^{G\times\tilde{P}_{r+1}}(\mathscr{Y}_{(r),(r)}). \tag{6.45}$$

Here Stab_T denotes the stabilizer of T in $G\times\tilde{P}_{r+1}$ and the notation $C_c^\infty - \mathrm{Ind}$ means that we are taking compactly supported induction. The isomorphism is given by the map $f\mapsto\hat{f}$, where we have set $\hat{f}(g,\tilde{p}) = [\omega_{(r),(r+1)}(g)\omega_{(r),(r+1)}(\tilde{p})f](T)$, for any given $f\in C_c^\infty(\mathcal{O})\otimes\mathscr{Y}_{(r),(r)}$. Its inverse is given by the map $h\mapsto\check{h}$, where $\check{h}(\tilde{p}^{-1}Tg) = \sigma_T(g)^{-1}\sigma_T(\tilde{p})^{-1}h(g,\tilde{p})$ for all $h\in C_c^\infty - \mathrm{Ind}_{\mathrm{Stab}_T}^{G\times\tilde{P}_{r+1}}(\mathscr{Y}_{(r),(r)})$, $g\in G$, $\tilde{p}\in\tilde{P}_{r+1}$. Let us describe the action on the right hand side of equation (6.45) more precisely. For this we will use the isomorphism $\mathscr{Y}_{(r),(r)}\cong\mathscr{S}_{-r,(r)}\otimes\mathscr{Y}_{(r-1),(r)}$. According to equations (6.22)–(6.28), the action of $\tilde{G}_{(r)}\tilde{N}_{r+1}$ on $\mathscr{Y}_{(r),(r)}$ is given by

$$\sigma_T(\tilde{g})h(S) = \omega_{(r-1),(r)}(\tilde{g})h(\tilde{g}^{-1}S) \qquad \text{for } \tilde{g}\in\tilde{G}_{(r)}, \tag{6.46}$$

$$\sigma_T(\exp\tilde{Z})h(S) = \psi(\mathrm{Tr}\,\tilde{Z}TT^*/2)h(S) \qquad \text{for } \tilde{Z}\in\tilde{\mathfrak{z}}_{r+1}, \tag{6.47}$$

$$\sigma_T(\exp\tilde{R})h(S) = \omega_{(r),(r)}(\tilde{R}T)h(S) = \psi(\mathrm{Tr}\,\tilde{R}TS^*)h(S) \qquad \text{for } \tilde{R}\in\mathrm{Hom}(\tilde{V}_{r+1},\tilde{V}_{(r)}). \tag{6.48}$$

In the last equality, we have used the fact that, since $T\in\mathrm{Hom}(V_r,\tilde{V}_{r+1})\subset\mathrm{Hom}(V_{(r)},\tilde{V}_{r+1})$, then $\tilde{R}T$ is in $\mathrm{Hom}(V_r,\tilde{V}_{(r)})\subset\mathrm{Hom}(V_{(r)},\tilde{V}_{(r)})$ which is a totally isotropic subspace that is complementary to $\mathrm{Hom}(V_{-r}, \tilde{V}_{(r)})$. (Here is where the choice of T simplifies matters.) Now observe that there is a continuous $G \times \tilde{P}_{r+1}$-intertwining map from $C_c^\infty(G \times \tilde{P}_{r+1} \times \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)}); \mathscr{Y}_{(r-1),(r)})$ to $C_c^\infty - \mathrm{Ind}_{\mathrm{Stab}_T}^{G\times\tilde{P}_{r+1}}(\mathscr{S}_{-r,(r)} \otimes \mathscr{Y}_{(r-1),(r)})$, with dense image, given by

$$\tilde{h}(g, \tilde{p}, S) = \int_{\mathrm{Stab}_T} (\sigma_T(x)^{-1} h(x_g g, x_{\tilde{p}}\tilde{p})(S)\, dx, \qquad h \in C_c^\infty(G \times \tilde{P}_{r+1} \times \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)}); \mathscr{Y}_{(r-1),(r)}).$$

Here x_g and $x_{\tilde{p}}$ are the projections of $x \in \mathrm{Stab}_T$ to G and $\tilde{P}_{r+1}$, respectively. Therefore, we may define a $\tilde{U}_{r+1}$-equivariant $\mathscr{Y}'_{(r-1),(r)}$-valued distribution $\bar{\lambda}$ on $G \times \tilde{P}_{r+1} \times \mathrm{Hom}(V_{-r}, V_{(r)})$ by setting

$$\bar{\lambda}(h) = \lambda(\check{\tilde{h}}), \qquad \text{for all } h \in C_c^\infty(G \times \tilde{P}_{r+1} \times \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)}); \mathscr{Y}_{(r-1),(r)}).$$

It is clear that if $\tilde{p} = \tilde{m}\tilde{g}\tilde{n}$, with $\tilde{m} \in \tilde{M}_{r+1} \cong \mathrm{GL}(\tilde{V}_{r+1})$, $\tilde{n} \in \tilde{N}_{r+1}$, and $\tilde{g} \in \tilde{G}_{(r)}$, then, for all $h \in C_c^\infty - \mathrm{Ind}_{\mathrm{Stab}_T}^{G\times\tilde{P}_{r+1}}(\mathscr{S}_{-r,(r)} \otimes \mathscr{Y}_{(r-1),(r)})$, $S \in \mathrm{Hom}(V_{-r}, \tilde{V}_{(r)})$, and $\tilde{R} \in \mathrm{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)})$, we have

$$\exp\tilde{R} \cdot h(g, \tilde{m}\tilde{g}\tilde{n}, S) = \psi(\mathrm{Tr}\,\tilde{R}\tilde{m}^{-1}TS^*\tilde{g}) h(g, \tilde{m}\tilde{g}\tilde{n}, S).$$

Hence

$$\exp\tilde{R} \cdot \bar{\lambda} = \psi(\mathrm{Tr}\,\tilde{R}\tilde{m}^{-1}TS^*\tilde{g})\bar{\lambda}.$$

In other words, the action of $\exp\tilde{R}$ on $\bar{\lambda}$ is given by multiplication by the function $(g, \tilde{m}\tilde{g}\tilde{n}, S) \mapsto \psi(\mathrm{Tr}\,\tilde{R}\tilde{m}^{-1}TS^*\tilde{g})$. On the other hand, since λ is $\tilde{U}_{r+1}$-equivariant, we should have

$$\exp\tilde{R} \cdot \bar{\lambda} = \psi(\mathrm{Tr}\,\tilde{R}\tilde{X})\bar{\lambda}, \qquad \text{for all } \tilde{R} \in \mathrm{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r-1)} \oplus \tilde{V}_{-r}) \subset \tilde{\mathfrak{u}}_{r+1}.$$

Here for the trace function, we are considering $\tilde{X}$ and $\tilde{R}$ as elements of $\mathrm{End}(\tilde{V}_{(r+1)})$. Therefore,

$$\mathrm{supp}\,\bar{\lambda} \subset \{(g, \tilde{m}\tilde{g}\tilde{n}, S) \mid \tilde{R}\tilde{m}^{-1}TS^*\tilde{g} = \tilde{R}\tilde{X}, \text{ for all } \tilde{R} \in \mathrm{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r-1)} \oplus \tilde{V}_{-r})\},$$

but the latter set is empty since $\mathrm{Im}\,\tilde{X}$ contains $\tilde{V}_{r+1}$, whereas the image of $\tilde{m}^{-1}TS^*\tilde{g}$ does not have the same property due to the fact that T is singular. From all this, we conclude that if $\mathcal{O} \subset \mathcal{A}$, then $D'(\mathcal{O}; \mathscr{Y}_{(r),(r)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} = 0$. This is statement (6.44) which, as we have already seen, implies statement (6.43).

Having proved statement (6.43), we come back to our original assertion that the map $\mathscr{Y}_{(r),(r+1)} \twoheadrightarrow (\mathscr{Y}_{(r),(r+1)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ factors through $\mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)}$. We have seen that any $\lambda \in (\mathscr{Y}'_{(r),(r+1)})^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ is completely determined by its restriction to $C_c^\infty(\mathcal{A}^c)$, where $\mathcal{A}^c$ is the complement of $\mathcal{A}$ in $\mathrm{Hom}(V_{(r)}, \tilde{V}_{r+1})$. But on $\mathcal{A}^c$, 0 is a regular value of the map $T \mapsto TT^*$. Therefore, $\lambda|_{A^c}$ is a distribution that *lives* on $\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})$, that is, λ can be identified with an element of $(\mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)})'$, as we wanted to show.

Given $f \in \mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)}$, we set

$$\tilde{f}_r(g) = [\omega_{(r),(r+1)}(g)f](T_r) = \omega_{(r),(r)}(g)[f(T_r g)], \qquad \text{for } g \in G. \tag{6.49}$$

Then it is clear that $\tilde{f}_r \in C^\infty(N_r G_{(r-1)} \backslash G; \mathscr{Y}_{(r),(r)})$ for all $f \in \mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)}$. Let

$$\mathscr{S}^+(N_r G_{(r-1)} \backslash G; \mathscr{Y}_{(r),(r)}) = \{\tilde{f}_r \mid f \in \mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)}\}.$$

Now observe that $\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})$ is a single orbit under the action of G. Therefore, since $G_{(r-1)}N_r$ is the stabilizer of $T_r \in \mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})$, we have that as G-modules

$$\mathscr{S}(\mathrm{Hom}_{GNM}(V_{(r)}, \tilde{V}_{r+1})) \otimes \mathscr{Y}_{(r),(r)} \cong \mathscr{S}^+(N_r G_{(r-1)} \backslash G; \mathscr{Y}_{(r),(r)}).$$

Now we will show that

$$\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{Y}_{(r),(r)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} \cong \mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{S}_{-r,-r}\otimes\mathscr{Y}_{(r-1),(r)}), \tag{6.50}$$

where

$$\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{S}_{-r,-r}\otimes\mathscr{Y}_{(r-1),(r)}) = \{f_r \mid f\in\mathscr{Y}_{(r),(r+1)}\}.$$

So let $\lambda\in(\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{Y}_{(r),(r)})')^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$. We will identify λ with a $\mathscr{Y}'_{(r-1),(r)}$-valued distribution on $G\times\mathrm{Hom}(V_{-r},\tilde{V}_{(r)})$. Observe that, for all $f\in\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{Y}_{(r),(r)})$, $g\in G$, $S\in\mathrm{Hom}(V_{-r},\tilde{V}_{(r)})$, and $\tilde{R}\in\mathrm{Hom}(\tilde{V}_{r+1},\tilde{V}_{(r)})$, we have

$$\exp\tilde{R}\cdot f(g)(S) = \psi(\mathrm{Tr}\,\tilde{R}T_rS^*)f(g)(S).$$

Therefore, since $\lambda\in(\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{Y}_{(r),(r)})')^{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$, we must have

$$\psi(\mathrm{Tr}\,\tilde{R}T_rS^*)\lambda = \psi(\mathrm{Tr}\,\tilde{R}\tilde{X})\lambda,$$

for all $\tilde{R}\in\mathrm{Hom}(\tilde{V}_{r+1},\tilde{V}_{(r-1)}\oplus\tilde{V}_{-r})\subset\tilde{\mathfrak{u}}_{r+1}$. Now, since $\tilde{X}$ is a regular value of the map $S\mapsto T_rS^*$, we conclude that λ is a distribution that *lives* on

$$G\times\{S\in\mathrm{Hom}(V_{-r},\tilde{V}_{(r)}) \mid \tilde{R}T_rS^* = \tilde{R}\tilde{X} \text{ for all } \tilde{R}\in\mathrm{Hom}(\tilde{V}_{r+1},\tilde{V}_{(r-1)}\oplus\tilde{V}_{-r})\}.$$

But the last set is just $G\times[T_{-r}+\mathrm{Hom}(V_{-r},\tilde{V}_{-r})]$. The isomorphism given in equation (6.50) now follows immediately.

The only thing that remains to be checked is that

$$\mathscr{S}^+(N_rG_{(r-1)}\backslash G;\mathscr{S}_{-r,-r}\otimes\mathscr{Y}_{(r-1),(r)}) \cong \mathscr{C}(N_rG_{(r-1)}\backslash G;\mathscr{S}_{-r,-r}\otimes\mathscr{Y}_{(r-1),(r)}). \tag{6.51}$$

For this, let ρ be a seminorm in $\mathscr{Y}_{(r-1),(r)}$, $Z_1\in\mathscr{D}(\mathrm{Hom}(V_{-r},\tilde{V}_{-r}))$, $Z_2\in U(\mathfrak{g})$, d_1, $d_2\in\mathbb{N}$ and $f\in\mathscr{S}_{(r),r+1}\otimes\mathscr{S}_{-r,(r)}\otimes\mathscr{Y}_{(r-1),(r)}$. Then

$$\begin{aligned}
q_{Z_1,Z_2,d_1,d_2,\rho}(f_r) &= \sup_{k\in K,m\in M_r,S\in\mathrm{Hom}(V_{-r},\tilde{V}_{-r})} \rho(Z_1[R_{Z_2}f_r(mk)](S))\|mk\|^{d_1}(1+\|S\|)^{d_2} \\
&= \sup_{k,m,S}\rho(Z_1[(k\cdot R_{Z_2}f)_r(m)](S))\|m\|^{d_1}(1+\|S\|)^{d_2} \\
&= \sup_{k,m,S}\rho(\omega_{(r),(r)}(m)Z_1(k\cdot R_{Z_2}f)(T_rm)(T_{-r}+S))\|m\|^{d_1}(1+\|S\|)^{d_2} \\
&= \sup_{k,m,S}\rho(Z_1(k\cdot R_{Z_2}f)(T_rm)(T_{-r}m+Sm))\|m\|^{d_1}(1+\|S\|)^{d_2}. \\
&\le \sup_{k,m,S} q_{Z_1,c_1,c_2,\rho}(k\cdot R_{Z_2}f)(1+\|T_rm+T_{-r}m\|)^{-c_1} \\
&\qquad \cdot(1+\|Sm\|)^{-c_2}\|m\|^{d_1}(1+\|S\|)^{d_2},
\end{aligned}$$

where c_1 ,$c_2\in\mathbb{N}$ are arbitrary and $q_{Z_1,c_1,c_2,\rho}$ is a seminorm on $\mathscr{S}_{(r),r+1}\otimes\mathscr{S}_{-r,(r)}\otimes\mathscr{Y}_{(r-1),(r)}$. But then there exists a constant $C>0$ such that

$$\begin{aligned}
q_{Z_1,Z_2,d_1,d_2,\rho}(f_r) &\le \sup_{k,m,S} C\,q_{Z_1,c_1,c_2,\rho}(k\cdot R_{Z_2}f)\|m\|^{-c_1}(1+\|S\|)^{-c_2}\|m\|^{c_2}\|m\|^{d_1}(1+\|S\|)^{d_2} \\
&\le \sup_{k,m,S} C\,q_{Z_1,c_1,c_2,\rho}(k\cdot R_{Z_2}f)\|m\|^{d_1+c_2-c_1}(1+\|S\|)^{d_2-c_2}.
\end{aligned}$$

Now observe that if we fix $c_2\gg d_2$ and $c_1\gg d_1+c_2$, then

$$\sup_{m,S} Cq_{Z_1,c_1,c_2,\rho}(k\cdot R_{Z_2}f)\|m\|^{d_1+c_2-c_1}(1+\|S\|)^{d_2-c_2}<\infty,$$

for all $k \in K$. Furthermore, the map $k \mapsto \sup_{m,S} C\, q_{Z_1,c_1,c_2,\rho}(k \cdot f)\|m\|^{d_1+c_2-c_1}(1+\|S\|)^{d_2-c_2}$ is continuous, and hence, since K is compact, bounded. Therefore $q_{Z_1,Z_2,d_1,d_2,\rho}(f_r) < \infty$ and (6.51) follows. □

Observe that the space $(\mathscr{Y}_{(r),(r+1)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}}$ carries a natural action of the group $\tilde{N}$. As a consequence of the isomorphism $(\mathscr{Y}_{(r),(r+1)})_{\tilde{U}_{r+1},\tilde{\chi}_{r+1}} \cong \mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$, we may define an action of $\tilde{N}$ on $\mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$ by the formula $(\tilde{n} \cdot f_r)(g)(S) := (\omega_{(r),(r+1)}(\tilde{n})f)_r(g)(S)$, for $f \in \mathscr{Y}_{(r),(r+1)}$, $\tilde{n} \in \tilde{N}$ and $S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{-r})$. We describe this action more concretely in the following

Lemma 6.6. *There is an induced action of* $\tilde{N}$ *(*$= \tilde{N}_{(r)}\tilde{N}_{r+1}$*) on* $\mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$*, as follows: for all* $\phi \in \mathscr{C}(N_r G_{(r-1)}\backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)})$*,* $g \in G$*,* $S \in \operatorname{Hom}(V_{-r}, \tilde{V}_{-r})$*,*

$$\begin{aligned}
(\exp \tilde{Z} \cdot \phi)(g)(S) &= \phi(g)(S) \qquad \text{for } \tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}\\
(\exp \tilde{R} \cdot \phi)(g)(S) &= \psi(\operatorname{Tr} \tilde{R} T_r T_{-r}^*)\psi(\operatorname{Tr} \tilde{R} T_r S^*)\phi(g)(S) \qquad \text{for } \tilde{R} \in \operatorname{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)})\\
(\exp \tilde{Z} \cdot \phi)(g)(S) &= \omega_{(r-1),(r)}(\exp \tilde{Z})[\phi(g)(S)] \qquad \text{for } \tilde{Z} \in \tilde{\mathfrak{z}}_r\\
(\exp \tilde{R} \cdot \phi)(g)(S) &= \omega_{(r-1),(r)}(\exp \tilde{R})[\phi(g)(S + \tilde{R}^* T_{-r})] \qquad \text{for } \tilde{R} \in \operatorname{Hom}(\tilde{V}_r, \tilde{V}_{(r-1)})\\
(\tilde{n} \cdot \phi)(g)(S) &= \omega_{(r-1),(r)}(\tilde{n})[\phi(g)(S)] \qquad \text{for } \tilde{n} \in \tilde{N}_{(r-1)}.
\end{aligned}$$

In particular, we have

$$(\tilde{u} \cdot \phi)(g)(S) = \omega_{(r-1),(r)}(\tilde{u})[\phi(g)(S)] \qquad \text{for } \tilde{u} \in \tilde{U}_{(r)}. \tag{6.52}$$

Proof. If $\tilde{Z} \in \tilde{\mathfrak{z}}_{r+1}$, we have already seen in equation (6.30) that

$$(\omega_{(r),(r+1)}(\exp \tilde{Z})f)_r(g)(S) = f_r(g)(S),$$

while if $\tilde{R} \in \operatorname{Hom}(\tilde{V}_{r+1}, \tilde{V}_{(r)}) \subset \tilde{\mathfrak{n}}_{r+1}$, similar to equation (6.31), we have

$$\begin{aligned}
(\omega_{(r),(r+1)}(\exp \tilde{R})f)_r(g)(S) &= \psi(\operatorname{Tr} \tilde{R} T_r (T_{-r} + S)^*) f_r(g)(S)\\
&= \psi(\operatorname{Tr} \tilde{R} T_r T_{-r}^*)\psi(\operatorname{Tr} \tilde{R} T_r S^*) f_r(g)(S).
\end{aligned}$$

On the other hand, if $\tilde{Z} \in \tilde{\mathfrak{z}}_r$, then

$$\begin{aligned}
(\omega_{(r),(r+1)}(\exp \tilde{Z})f)_r(g)(S) &= (\omega_{(r),(r+1)}(\exp \tilde{Z})[\omega_{(r),(r+1)}(g)f])(T_r, T_{-r} + S)\\
&= (\omega_{(r),(r)}(\exp \tilde{Z})[(\omega_{(r),(r+1)}(g)f)(T_r)])(T_{-r} + S)\\
&= \omega_{(r-1),(r)}(\exp \tilde{Z})([\omega_{(r),(r+1)}(g)f](T_r, (\exp \tilde{Z})^{-1}(T_{-r} + S)))\\
&= \omega_{(r-1),(r)}(\exp \tilde{Z})([\omega_{(r),(r+1)}(g)f](T_r, T_{-r} + S))\\
&= \omega_{(r-1),(r)}(\exp \tilde{Z}) f_r(g)(S).
\end{aligned}$$

Here we are using that $\tilde{Z}(T_{-r} + S) = 0$ because $\operatorname{Im}(T_{-r} + S) \subset \tilde{V}_{-r+1} \oplus \tilde{V}_{-r}$ and $\tilde{Z}|_{\tilde{V}_{-r+1} \oplus \tilde{V}_{-r}} = 0$. On the other hand, if $\tilde{R} \in \operatorname{Hom}(\tilde{V}_r, \tilde{V}_{(r-1)}) \subset \tilde{\mathfrak{n}}_r$, then, according to equations (6.46), (6.47) and (6.48)

$$\begin{aligned}
(\omega_{(r),(r+1)}(\exp \tilde{R})f)_r(g)(S) &= (\omega_{(r),(r+1)}(\exp \tilde{R})[\omega_{(r),(r+1)}(g)f])(T_r, T_{-r} + S)\\
&= (\omega_{(r),(r)}(\exp \tilde{R})[(\omega_{(r),(r+1)}(g)f)(T_r)])(T_{-r} + S)\\
&= \omega_{(r-1),(r)}(\exp \tilde{R})([\omega_{(r),(r+1)}(g)f](T_r, (\exp \tilde{R})^{-1}(T_{-r} + S)))\\
&= \omega_{(r-1),(r)}(\exp \tilde{R})([\omega_{(r),(r+1)}(g)f](T_r, T_{-r} + S + \tilde{R}^* T_{-r}))\\
&= \omega_{(r-1),(r)}(\exp \tilde{R}) f_r(g)(S + \tilde{R}^* T_{-r}).
\end{aligned}$$

Here we are using the identification $\mathrm{Hom}(\tilde V_r, \tilde V_{(r-1)}) \ni \tilde R \leftrightarrow \tilde R - \tilde R^* \in \tilde{\mathfrak{n}}_r$ as in equation (6.20). We are also using that, since $T_{-r} \in \mathrm{Hom}(V_{-r}, \tilde V_{-r+1})$, then $\tilde R^* T_{-r} \in \mathrm{Hom}(V_{-r}, \tilde V_{-r})$ and that

$$\tilde R^* S_{-r} = \tilde R S_{-r} = \tilde R T_{-r} = \tilde R^* \tilde R T_{-r} = \tilde R^* \tilde R S_{-r} = 0,$$

for all $\tilde R \in \mathrm{Hom}(\tilde V_r, \tilde V_{(r-1)})$, $S \in \mathrm{Hom}(V_{-r}, \tilde V_{-r})$. We note, in particular, that $\tilde R^* T_{-r} = 0$ if $\tilde R \in \mathrm{Hom}(\tilde V_r, \tilde V_{(r-2)} \oplus V_{-r+1}) \subset \tilde{\mathfrak{u}}_r$. Finally, if $\tilde n \in \tilde N_{(r-1)} \subset \tilde G_{(r)}$, then

$$\begin{aligned}
(\omega_{(r),(r+1)}(\tilde n) f)_r(g)(S) &= (\omega_{(r),(r+1)}(\tilde n)[\omega_{(r),(r+1)}(g) f])(T_r, T_{-r} + S) \\
&= (\omega_{(r),(r)}(\tilde n)[(\omega_{(r),(r+1)}(g) f)(T_r)])(T_{-r} + S) \\
&= \omega_{(r-1),(r)}(\tilde n)([\omega_{(r),(r+1)}(g) f](T_r, \tilde n^{-1}(T_{-r} + S))) \\
&= \omega_{(r-1),(r)}(\tilde n)([\omega_{(r),(r+1)}(g) f](T_r, T_{-r} + S)) \\
&= \omega_{(r-1),(r)}(\tilde n) f_r(g)(S).
\end{aligned}$$

Here we are using that, considered as an element of $\tilde G_{(r)}$, $\tilde n \in \tilde N_{(r-1)}$ leaves $\tilde V_{-r+1} \oplus \tilde V_{-r}$ fixed. ☐

6.5. **From $\mathscr{Y}_{(r),(r+1)}$ to $\mathscr{H}_{\gamma,\tilde\gamma}$: proof of Proposition 6.4.** We are finally ready to complete the proof of the key proposition, to which we refer the reader for the unexplained notation below.

Proof (of Proposition 6.4). We will prove the result by induction on r. The case $r = 0$ is immediate. Observe that $\tilde U = \tilde U_{(r)} \tilde U_{r+1}$ and $\tilde\chi_{(r+1)} = \tilde\chi_{(r)} \tilde\chi_{r+1}$. Therefore, from Lemma 6.5

$$\begin{aligned}
(\mathscr{Y}_{(r),(r+1)})_{\tilde U, \tilde\chi} &\cong (\mathscr{C}(N_r G_{(r-1)} \backslash G; \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}))_{\tilde U_{(r)}, \tilde\chi_{(r)}} \\
&\cong \mathscr{C}(N_r G_{(r-1)} \backslash G; \mathscr{S}_{-r,-r} \otimes (\mathscr{Y}_{(r-1),(r)})_{\tilde U_{(r)}, \tilde\chi_{(r)}}),
\end{aligned}$$

where the last equation follows from the fact that $\tilde U_{(r)}$ acts only on the values. See equation (6.52). Now, given $f \in \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}$, $g \in G$ and $S_{-k} \in \mathrm{Hom}(V_{-k}, \tilde V_{-k})$, $k = 1, \ldots, l$, set

$$\begin{aligned}
f_{(r-1)}(g)(S_{-r}, \ldots, S_{-1}) &= (\bar\sigma_{T_r}(g) f)(S_{-r}, T_{r-1}, T_{-r+1} + S_{-r+1}, \ldots, T_0) \\
&= (\omega_{(r-1),(r)}(g)[f(S_{-r})])(T_{r-1}, T_{-r+1} + S_{-r+1}, \ldots, T_0).
\end{aligned}$$

For convenience, we denote the representation $\tau^T_{\gamma,\tilde\gamma}$ on $\mathscr{H}_{\gamma,\tilde\gamma}$ (defined in equation (6.7)) by the more concise symbol $\mathscr{H}_T$. Then, by induction hypothesis, the map $f \mapsto f_{(r-1)}$ induces a $G_{(r-1)}$-intertwining isomorphism

$$\Psi_{(r-1)} : \mathscr{S}_{-r,-r} \otimes (\mathscr{Y}_{(r-1),(r)})_{\tilde U_{(r)}, \tilde\chi_{(r)}} \longrightarrow \mathscr{C}(N_{(r-1)} \backslash G_{(r-1)}; \mathscr{H}_{T_{(r)}}),$$

where $T_{(l)} = \oplus_{k=-l}^{l} T_k$, for $l = 0, \ldots, r$. Notice that $T_{(r)} = T_{\gamma,\tilde\gamma}$. Also notice that in the above equation we are identifying the spaces $\mathscr{S}_{-r,-r} \otimes \mathscr{C}(N_{(r-1)} \backslash G_{(r-1)}; \mathscr{H}_{T_{(r-1)}}) \cong \mathscr{C}(N_{(r-1)} \backslash G_{(r-1)}; \mathscr{H}_{T_{(r)}})$. We claim that $\Psi_{(r-1)}$ is actually an isomorphism of $G_{(r-1)} N_r$-modules:

$$\Psi_{(r-1)} : \mathscr{S}_{-r,-r} \otimes (\mathscr{Y}_{(r-1),(r)})_{\tilde U_{(r)}, \tilde\chi_{(r)}} \longrightarrow \mathscr{C}(N \backslash G_{(r-1)} N_r; \mathscr{H}_{T_{(r)}}), \tag{6.53}$$

where

$$\mathscr{C}(N \backslash G_{(r-1)} N_r; \mathscr{H}_{T_{(r)}}) = \{ f \in C^\infty(N \backslash G_{(r-1)} N_r; \mathscr{H}_{T_{(r)}}) \mid f|_{G_{(r-1)}} \in \mathscr{C}(N_{(r-1)} \backslash G_{(r-1)}; \mathscr{H}_{T_{(r-1)}}).\}$$

Observe that if $f \in \mathscr{C}(N \backslash G_{(r-1)} N_r; \mathscr{H}_{T_{(r)}})$, $g \in G_{(r-1)}$ and $n \in N_r$, then

$$(n \cdot f)(g) = f(gn) = f((gng^{-1})g) = \tau_{T_{(r)}}(gng^{-1}) f(g).$$

Therefore, to prove our claim, we need to show that for all $f \in \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}$, $g \in G_{(r-1)}$, $n \in N_r$,

$$(\bar\sigma_{T_r}(n) f)_{(r-1)}(g) = \tau_{T_{(r)}}(gng^{-1}) f_{(r-1)}(g). \tag{6.54}$$

Now, by definition,

$$\begin{aligned}(\bar{\sigma}_{T_r}(n)f)_{(r-1)}(g)(S_{-r},\dots,S_{-1}) &= (\bar{\sigma}_{T_r}(g)\bar{\sigma}_{T_r}(n)f)(S_{-r},T_{r-1},T_{-r+1}+S_{-r+1},\dots,T_0)\\ &= (\bar{\sigma}_{T_r}(gng^{-1})[\bar{\sigma}_{T_r}(g)f])(S_{-r},T_{r-1},T_{-r+1}+S_{-r+1},\dots,T_0)\\ &= (\bar{\sigma}_{T_r}(gng^{-1})[\bar{\sigma}_{T_r}(g)f])_{(r-1)}(e)(S_{-r},S_{-r+1},\dots,S_{-1}).\end{aligned}$$

From all this we see that in order to prove the claim given in equation (6.53) it suffices to show that for all $f \in \mathscr{S}_{-r,-r} \otimes \mathscr{Y}_{(r-1),(r)}$, $n \in N_r$,

$$(\bar{\sigma}_{T_r}(n)f)_{(r-1)}(e) = \tau_{T_{(r)}}(n)f_{(r-1)}(e). \tag{6.55}$$

Now, by equation (6.35), if $Z \in \mathfrak{z}_r$, then

$$(\bar{\sigma}_{T_r}(\exp Z)f)_{(r-1)}(e)(S_{-r},\dots,S_{-1}) = f(S_{-r},T_{r-1},\dots,T_0) = f_{(r-1)}(e)(S_{-r},\dots,S_{-1}).$$

On the other hand, by equation (6.36), if $R \in \operatorname{Hom}(V_{(r-2)} \oplus V_{r-1}, V_{-r}) \subset \mathfrak{u}_r$, then

$$\begin{aligned}(\bar{\sigma}_{T_r}(\exp R)f)_{(r-1)}(e)(S_{-r},\dots,S_{-1}) &= (\omega_{(r-1),(r)}(T_{-r}R)[f(S_{-r})])(T_{r-1},\dots,T_0)\\ &= \psi(-\operatorname{Tr} T_{-r}RT^*_{r-2})f(S_{-r},T_{r-1},\dots,T_0)\\ &= \chi_\gamma(\exp R)^{-1}f_{(r-1)}(e)(S_{-r},\dots,S_{-1}),\end{aligned}$$

in other words, for all $u \in U_r$,

$$(\bar{\sigma}_{T_r}(u)f)_{(r-1)}(e)(S_{-r},\dots,S_{-1}) = \chi_\gamma(u)^{-1}f_{(r-1)}(e)(S_{-r},\dots,S_{-1}).$$

Finally, again by equation (6.36), if $R \in \mathfrak{g}_{-1} \cap \mathfrak{n}_r \cong \operatorname{Hom}(V_{-r+1}, V_{-r})$, then

$$\begin{aligned}(\bar{\sigma}_{T_r}(\exp R)f)_{(r-1)}(e)(S_{-r},\dots,S_{-1}) &= (\omega_{(r-1),(r)}(T_{-r}R+S_{-r}R)[f(S_{-r})](T_{r-1},\dots,T_0)\\ &= \psi(-\operatorname{Tr} S_{-r}RT^*_{r-1})f_{(r-1)}(e)(S_{-r},S_{-r+1}+T_{-r}R,\dots,S_{-1}).\end{aligned}$$

Now, for such an R, $\sigma_{T_{(r)}}(R,0) = T_{-r}R - T_{r-1}R^*$. Therefore,

$$\begin{aligned}\tau_{T_{(r)}}(\exp R)f_{(r-1)}(e)(S_{-r},\dots,S_{-1}) &= \tau_{\gamma,\tilde{\gamma}}(T_{-r}R - T_{r-1}R^*)f_{(r-1)}(e)(S_{-r},\dots,S_{-1})\\ &= \psi(-\operatorname{Tr} S_{-r}RT^*_{r-1})f_{(r-1)}(e)(S_{-r},S_{-r+1}+T_{-r}R,\dots,S_{-1}).\end{aligned}$$

Therefore, equation (6.55) holds true, and hence we have proved our claim given in equation (6.53).

We will now show that

$$\mathscr{C}(N\backslash G;\mathscr{H}_{T_{(r)}}) \cong \mathscr{C}(N_rG_{(r-1)}\backslash G;\mathscr{C}(N\backslash G_{(r-1)}N_r;\mathscr{H}_{T_{(r)}})).$$

Given $f \in \mathscr{C}(N_rG_{(r-1)}\backslash G;\mathscr{C}(N\backslash G_{(r-1)}N_r;\mathscr{H}_{T_{(r)}}))$, set $\check{f}(g) = f(g)(e)$, for all $g \in G$. Then it is clear that $\check{f} \in C^\infty(N\backslash G;\mathscr{H}_{T_{(r)}})$, but we claim that $\check{f}$ is actually in $\mathscr{C}(N\backslash G;\mathscr{H}_{T_{(r)}})$. Effectively, given a semi-norm ρ of $\mathscr{H}_{T_{(r)}}$, $Z \in U(\mathfrak{g})$ and $d \in \mathbb{N}$,

$$\begin{aligned}q_{Z,d,\rho}(\check{f}) &= \sup_{k\in K,\, m_1\in M_r,\, m_2\in G_{(r-1)}} \rho(R_Z\check{f}(m_2mk_1))\|m_1m_2\|^d\\ &= \sup_{k,m_1,m_2} \rho(R_Zf(m_2mk_1)(e))\|m_1m_2\|^d\\ &\le \sup_{k,m_1,m_2} \rho(R_Zf(mk_1)(m_2^{-1}))\|m_1\|^d\|m_2\|^d\\ &= p_{Z,1,d,d,\rho}(f) < \infty.\end{aligned}$$

Now given $f \in \mathscr{C}(N\backslash G;\mathscr{H}_{T_{(r)}})$, set $\hat{f}(g)(h) = f(hg)$. Then again it is clear that

$$\hat{f} \in C^\infty(N_rG_{(r-1)}\backslash G;C^\infty(N\backslash N_rG_{(r-1)};\mathscr{H}_{T_{(r)}})),$$

but we claim that, actually, $\hat{f}$ is a function in $\mathscr{C}(N_rG_{(r-1)}\backslash G;\mathscr{C}(N\backslash G_{(r-1)}N_r;\mathscr{H}_{T_{(r)}}))$. Effectively, let ρ be a semi-norm of $\mathscr{H}_{T_{(r)}}$, $Z_1 \in U(\mathfrak{g})$, $Z_2 \in U(\mathfrak{g}_{(r-1)})$ and $d_1, d_2 \in \mathbb{N}$. Then $Z_2 \in U^m(\mathfrak{g})$ for some

m and hence there exists a basis $\{Y_j\}_{j=1}^{l}$ of $U^m(\mathfrak{g})$ such that for all $g \in G$, $\mathrm{Ad}(g)Z_2 = \sum_j a_j(g)Y_j$ for some functions a_j. Since $(\mathrm{Ad}, U^m(\mathfrak{g}))$ is a finite dimensional representation, there exists a constant d_m such that $|a(g)| \leq \|g\|^{d_m}$ for all $g \in G$. Taking this into account we have that

$$\begin{aligned} p_{Z_1,Z_2,d_1,d_2,\rho}(\hat{f}) &= \sup_{k\in K,\, m_1\in M_r,\, m_2\in G_{(r-1)}} \rho(R_{Z_2}(R_{Z_1}\hat{f}(mk_1))(m_2))\|m_1\|^{d_1}\|m_2\|^{d_2} \\ &= \sup_{k,m_1,m_2} \rho(R_{\mathrm{Ad}(mk_1)^{-1}Z_2}R_{Z_1}f(m_2mk_1))\|m_1\|^{d_1}\|m_2\|^{d_2} \\ &\leq \sup_{k,m_1,m_2} \sum_{j=1}^{l} \rho(a(mk_1)R_{Y_j}R_{Z_1}f(mk_1m_2))\|m_1\|^{d_1}\|m_2\|^{d_2} \\ &\leq \sup_{k,m_1,m_2} \sum_{j=1}^{l} \rho(R_{Y_j}R_{Z_1}f(mk_1m_2))\|m_1\|^{d_1+d_m}\|m_2\|^{d_2} \\ &\leq \sup_{k,m_1,m_2} \sum_{j=1}^{l} \rho(R_{Y_jZ_1}f(mk_1m_2))\|m_1m_2\|^{d_1+d_m+d_2} \\ &= \sum_{j=1}^{l} q_{Y_jZ_1,d_1+d_2+d_m,\rho}(f) < \infty. \end{aligned}$$

Here we have used the fact that $\|m_1m_2\| = \max\{\|m_1\|, \|m_2\|\} \geq \|m_1\|, \|m_2\|$. Now the only thing left to be check is that $\hat{\check{f}} = f$ and $\check{\hat{f}} = f$, but by definition

$$\hat{\check{f}}(g)(h) = \check{f}(gh) = f(gh)(e) = f(g)(h),$$

and similarly $\check{\hat{f}} = f$. $\square$

Appendix A. Some facts on vector valued distributions

In this appendix, we collect some facts about vector valued distributions. For a more detailed presentation of the material, the interested reader should refer to [14], over which this appendix is based.

A.1. **Vector valued distributions.** Let $\mathscr{X}$ be a C^∞-manifold and $\mathscr{V}$ a Fréchet space. We write $C^\infty(\mathscr{X};\mathscr{V})$ and $C_c^\infty(\mathscr{X};\mathscr{V})$ for the topological vector spaces of the C^∞ mappings from $\mathscr{X}$ to $\mathscr{V}$, and those with compact support, respectively. Let $\mathscr{V}'$ be the dual of $\mathscr{V}$ provided with the strong dual topology, that is, the topology of uniform convergence on bounded subsets of $\mathscr{V}$. Finally, let $\mathrm{Diff}(\mathscr{X})$ be the sheaf of differential operators on $\mathscr{X}$.

Definition. A $\mathscr{V}$-distribution on $\mathscr{X}$ is an element of the topological vector space

$$D'(\mathscr{X};\mathscr{V}) := C_c^\infty(\mathscr{X};\mathscr{V})' \cong (C_c^\infty(\mathscr{X}) \otimes \mathscr{V})'$$

consisting of the continuous linear forms on $C_c^\infty(\mathscr{X};\mathscr{V})$, provided with the strong dual topology. A $\mathscr{V}'$-valued distribution on $\mathscr{X}$ is an element of the space $\mathrm{Lin}(C_c^\infty(\mathscr{X});\mathscr{V}')$ of continuous linear maps from $C_c^\infty(\mathscr{X})$ to $\mathscr{V}'$. We will also equip $\mathrm{Lin}(C_c^\infty(\mathscr{X});\mathscr{V}')$ with the strong dual topology.

A $\mathscr{V}$ distribution λ defines canonically a $\mathscr{V}'$-valued distribution $\tilde{\lambda}$ on $\mathscr{X}$, that is, we have a canonical isomorphism

$$\lambda \longleftrightarrow \tilde{\lambda} \qquad D'(\mathscr{X};\mathscr{V}) \longleftrightarrow \mathrm{Lin}(C_c^\infty(\mathscr{X});\mathscr{V}').$$

In fact, if $\lambda \in D'(\mathscr{X};\mathscr{V})$ and $f \in C_c^\infty(\mathscr{X})$, then $\tilde{\lambda}(f) \in \mathscr{V}'$ is well-defined by $\tilde{\lambda}(f)(v) = \lambda(f \otimes v)$, for all $v \in \mathscr{V}$. In what follows we will frequently identify these two spaces.

A.2. **Transverse order of a $\mathscr{V}$-distribution.**

Definition. A $\mathscr{V}$-distribution $\lambda \in D'(\mathscr{X};\mathscr{V})$ is said to be of order $\leq l$ if for every compact set $\mathcal{K} \subset \mathscr{X}$ there exists a constant $c > 0$, $Z_1, \dots, Z_m \in \mathrm{Diff}^{(l)}(\mathscr{X})$ and a seminorm ν on $\mathscr{V}$ such that for all $f \in C_c^\infty(\mathscr{X};\mathscr{V})$

$$|\lambda(f)| \leq c \sum_{j=1}^{m} \sup_{x \in \mathcal{K}} \nu(Z_j f(x)).$$

We write $D'^{(l)}(\mathscr{X};\mathscr{V})$ for the subspace of $D'(\mathscr{X};\mathscr{V})$ consisting of elements of order $\leq l$.

Note that every $\mathscr{V}$-distribution on $\mathscr{X}$ is of finite order over every relatively compact open subset of $\mathscr{X}$.

Suppose $\mathcal{C}$ is a closed C^∞-manifold that is regularly embedded in $\mathscr{X}$. We write $D'_{\mathcal{C}}(\mathscr{X};\mathscr{V})$ for the distributions in $D'(\mathscr{X};\mathscr{V})$ with support contained in $\mathcal{C}$. We say that $\lambda \in D'(\mathscr{X};\mathscr{V})$ has transverse order $\leq l$ at $x \in \mathcal{C}$ if the following holds: there exists an open neighborhood U_x of x in $\mathscr{X}$ such that

$$\lambda(f) = 0, \qquad \begin{array}{c} \text{if } f \in C_c^\infty(U_x;\mathscr{V}) \text{ satisfies } Zf|_{\mathcal{C}\cap U_x} = 0 \\ \text{for all } Z \in \mathrm{Diff}^{(l)}(U_x). \end{array}$$

If λ satisfies this definition, then the transverse order of λ is $\leq l$ at all points of $\mathcal{C}$ in a suitable neighborhood of x. Let $D'^{(l)}_{\mathcal{C}}(\mathscr{X};\mathscr{V})$ be the linear subspace of elements in $D'(\mathscr{X};\mathscr{V})$ which have transverse order $\leq l$ for all points of $\mathcal{C}$. We say that λ *lives* on $\mathcal{C}$ if λ belongs to $D'(\mathcal{C};\mathscr{V}) := D'^{(0)}_{\mathcal{C}}(\mathscr{X};\mathscr{V})$. This is the same as saying that $\lambda(f) = 0$ if $f \in C_c^\infty(\mathscr{X};\mathscr{V})$ satisfies $f|_{\mathcal{C}} = 0$. In this case there exists a unique distribution μ on $\mathcal{C}$ such that $\lambda(f) = \mu(f|_{\mathcal{C}})$, for all $f \in C_c^\infty(\mathscr{X};\mathscr{V})$; and often we will identify λ with μ. Note that in this case $\lambda \in D'^{(r)}_{\mathcal{C}}(\mathscr{X};\mathscr{V})$ and $\mathrm{supp}(\lambda) \subset \mathcal{C}$.

A.3. **Transverse jet bundle.** As before, let $\mathscr{X}$ be a C^∞-manifold, and let $\mathcal{C} \subset \mathscr{X}$ be a closed C^∞-manifold regularly embedded in $\mathscr{X}$. For any $x \in \mathcal{C}$, let C_x^∞ be the algebra of germs of C^∞ functions defined around x and $\mathrm{Diff}_x^{(l)}$ the C_x^∞-module of germs of differential operators of order $\leq l$ defined around x. Let $V_x^{(l)}$ be the C_x^∞-submodule of $\mathrm{Diff}_x^{(l)}$ generated by germs of l-tuples $(v_1, \dots, v_l)$ of vector fields around x for which at least one of the v_i's is tangent to $\mathcal{C}$; let $I_x^{(l)} = \mathrm{Diff}_x^{(l-1)} + V_x^{(l)}$. By choosing local coordinates at x, it is easy to see that $I_x^{(l)}$ actually is the stalk at x of a subsheaf $I^{(l)}$ of the sheaf $\mathrm{Diff}^{(l)}$ of differential operators of order $\leq l$ on $\mathscr{X}$. Hence we have a well-defined quotient sheaf

$$M^{(l)} = \mathrm{Diff}^{(l)} / I^{(l)},$$

with stalk at x equal to $M_x^{(r)} = \mathrm{Diff}_x^{(r)} / I_x^{(r)}$. We write $\partial \mapsto \overline{\partial}$ for the projection $\mathrm{Diff}_x^{(r)} \longrightarrow M_x^{(r)}$. Since $\mathrm{Diff}_x^{(r)}$ and $I_x^{(r)}$ are both C_x^∞-modules, $M_x^{(r)}$ also has the structure of a C_x^∞-module, and the projection is a mapping of C_x^∞-modules.

Given $x \in \mathcal{C}$ we can select an open neighborhood U_x of X and local coordinates $(t : u) = (t_1, \dots, t_p, u_1, \dots, u_q)$ on U_x such that

$$U_x \cap \mathcal{C} = \{(t : u) \,|\, t_1 = \ldots = t_p = 0\}.$$

It is easy to show that the elements $\overline{\partial_t^\alpha}$ $(|\alpha| = l)$ form a free basis for the sections of $M^{(l)}$ around x. Here $\alpha = (\alpha_1, \dots, \alpha_p)$ is a multi-index, $|\alpha| = \alpha_1 + \cdots + \alpha_p$ and $\partial_t^\alpha = \partial^{|\alpha|}/\partial_{t_1}^{\alpha_1} \cdots \partial_{t_p}^{\alpha_p}$. Thus $M^{(l)}$

is a vector bundle over $\mathcal{C}$ of finite rank. This is the l-th graded part of the transverse jet bundle on $\mathcal{C}$. We have that $M^{(l)}$ is the l-th symmetric power of $M^{(1)}$. We write $M^{(l)'}$ for the dual to $M^{(l)}$.

A.4. **Transverse symbols.** In view of the isomorphism $M^{(l)} \otimes (C_c^\infty(\mathcal{C}) \otimes \mathscr{V})' \cong (C_c^\infty(\mathcal{C}) \otimes (M^{(l)'} \otimes \otimes \mathscr{V}))'$ it follows that

$$M^{(l)} \otimes D'(\mathcal{C}; \mathscr{V}) \cong D'(\mathcal{C}; M^{(l)'} \otimes \mathscr{V}).$$

Let $T \in D'^{(l)}_{\mathcal{C}}(\mathcal{X}; \mathscr{V})$. Given $x \in \mathcal{C}$ choose a neighborhood U_x of x and a system of local coordinates $(t : u)$ as before. Then there exist uniquely determined distributions $\tau_\alpha \in D'(\mathcal{C}; \mathscr{V})$ with $|\alpha| \leq l$, such that on U_x we have

$$T = \sum_{|\alpha| \leq l} (-1)^{|\alpha|} \tau_\alpha \partial_t^\alpha. \tag{A.1}$$

Theorem. *For any* $T \in D'^{(l)}_{\mathcal{C}}(\mathcal{X}; \mathscr{V})$ *there is a* $\sigma(T) = \sigma^{(l)}(T) \in D'(\mathcal{C}; M^{(l)'} \otimes \mathscr{V})$ *with the following property: if* T *has an expression in local coordinates as in equation (A.1), then*

$$\sigma(T) = \sum_{|\alpha| = l} \overline{\partial_t^\alpha} \otimes \tau_\alpha.$$

The element $\sigma(T)$ *is uniquely determined by this requirement. The map*

$$\sigma : D'^{(l)}_{\mathcal{C}}(\mathcal{X}; \mathscr{V}) \longrightarrow D'(\mathcal{C}; M^{(l)'} \otimes \mathscr{V})$$

is injective modulo the linear subspace $D'^{(l-1)}_{\mathcal{C}}(X; \mathscr{V})$ *and is independent of the choice of local coordinate systems.*

References

[1] D. Barbasch and D. A. Vogan Jr., *The local structure of characters,* J. Funct. Anal. 37, (1980), no. 1, 27–55.

[2] D. H. Collingwood and W. M. McGovern, *Nilpotent Orbits in Semisimple Lie Algebras,* Van Nostrand Reinhold Mathematics Series, 1992.

[3] A. Daszkiewicz, W. Kraskiewicz and T. Przebinda, *Nilpotent orbits and complex dual pairs,* J. Algebra 190, (1997), 518–539.

[4] F. du Cloux, *Sur les reprsentations diffrentiables des groupes de Lie algbriques*, Ann. Sci. Ecole Norm. Sup. 24, (1991), no. 3, 257–318.

[5] A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires,* Mem. Amer. Math. Soc. 1955, no. 16, 140 pp.

[6] Harish-Chandra, *Admissible invariant distributions on reductive p-adic groups,* Queen's Papers in Pure and Applied Math. 48, (1978), 281–347.

[7] R. Howe, *θ-series and invariant theory*, in Automorphic Forms, Representations and L-functions, Proc. Symp. Pure Math. 33, (1979), pp. 275–285.

[8] R. Howe, *L^2 duality for stable reductive dual pairs*, unpublished.

[9] R. Howe, *The oscillator representation: algebraic and analytic preliminaries*, unpublished notes.

[10] R. Howe, *Wave front sets of representations of Lie groups*, in Automorphic forms, representation theory and arithmetic (Bombay, 1979), pp. 117–140, Tata Inst. Fund. Res. Studies in Math., 10, Tata Inst. Fundamental Res., Bombay, 1981.

[11] R. Howe, *Transcending classical invariant theory*, J. Amer. Math. Soc. 2, (1989), 535–552.

[12] D. Jiang, *Periods of automorphic forms*, Proceedings of the International Conference on Complex Geometry and Related Fields, Studies in Advanced Mathematics, Vol 39, 2007, pp. 125–148, American Mathematical Society and International Press.

[13] N. Kawanaka, "Generalized Gelfand-Graev representations and Ennola duality", in *Algebraic groups and related topics,* Advanced Studies in Pure Math. 6, (1985), pp. 175–206.

[14] J. A. C. Kolk and V. S. Varadarajan, *On the transverse symbol of vectorial distributions and some applications to harmonic analysis,* Indag. Math. (N.S.) 7, (1996), no. 1, 67–96.

[15] B. Kostant, *On Whittaker vectors and representation theory,* Invent. math. 48, (1978), 101–184.

[16] H. Kraft and C. Procesi, *On the geometry of conjugate classes in classical groups,* Comment. Math. Helvetici 57, (1982), 539–602.
[17] S. S. Kuda, *On the local theta-correspondence*, Invent. math. 83, (1986), 229–255.
[18] J.-S. Li, *Singular unitary representations of classical groups,*, Invent. math. 97, (1989), 237–255.
[19] H.Y. Loke and J.-J. Ma, *Invariants and K-spectrums of local theta lifts*, preprint, (2013). arXiv:1302.1031.
[20] H. Matumoto, *$C^{-\infty}$-Whittaker vectors corresponding to a principal nilpotent orbit of a real reductive linear Lie group, and wave front sets,* Compositio Math. 82, (1992), 189–244.
[21] C. Mœglin, *Correspondance de Howe et front d'onde,* Adv. Math. 133, (1998), no. 2, 224–285.
[22] C. Mœglin, M.-F. Vigneras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps p-adique*, Lecture Notes in Mathematics, vol. 1291, Springer-Verlag, 1987.
[23] C. Mœglin and J.-L. Waldspurger, *Modèles de Whittaker dégénérés pour des groupes p-adiques,* Math. Z. 196, (1987), 427–452.
[24] K. Nishiyama, H. Ochiai and C.-B. Zhu, *Theta lifting of nilpotent orbits for symmetric pairs,* Trans. Amer. Math. Soc. 358, (2006), no. 6, 2713–2734.
[25] K. Nishiyama and C.-B. Zhu, *Theta lifting of unitary lowest weight modules and their associated cycles,* Duke Math. J. 125, (2004), no. 3, 415–465.
[26] M. E. Novodvorskii and I. Piatecki-Shapiro, *Generalized Bessel models for a symplectic group of rank* 2, (Russian) Mat. Sb. (N.S.) 90 (132), (1973), 246–256.
[27] S.-Y. Pan, *Orbit correspondences for real reductive pairs,* Pacific J. Math. 248, (2010), no. 2, 403–427.
[28] V. Protsak and T. Przebinda, *On the occurence of admissible representations in the real Howe correspondence in stable range*, Manuscripta Math. 126, (2008), 135–141.
[29] T. Przebinda, *Characters, dual pairs, and unipotent representations*, J. Funct. Anal. 98, (1991), no. 1, 59–96.
[30] T. Przebinda, *Characters, dual pairs, and unitary representations*, Duke Math. J. 69, (1993), no. 3, 547–592.
[31] R. Ranga Rao, *On some explicit formulas in the theory of Weil representation,* Pacific J. Math. 157, (1993), no. 2, 335–371.
[32] W. Schmid and K. Vilonen, *Characteristic cycles and wave front cycles of representations of reductive Lie groups,* Ann. of Math. 151, (2000), no. 3, 1071–1118.
[33] J. Sekiguchi, *Remarks on real nilpotent orbits of a symmetric pair,* J. Math. Soc. Japan 39, (1987), no. 1, 127–138.
[34] J. A. Shalika, *The multiplicity one theorem for* GL_n*,* Ann. of Math. 100, (1974), 171–193.
[35] B. Sun, *On representations of real Jacobi groups*, Sci. China Math. 55, (2012), 541–555.
[36] B. Sun and C.-B. Zhu, *A general form of Gelfand-Kazhdan criterion,* Manuscripta Math. 136, (2011), 185–197.
[37] D. A. Vogan Jr., *Gelfand-Kirillov dimension for Harish-Chandra modules,* Invent. math. 48, (1978), 75–98.
[38] D. A. Vogan Jr., "Associated varieties and unipotent representations" in *Harmonic analysis on reductive groups* (Brunswick, ME, 1989), pp. 315–388, Progr. Math., 101, Birkhäuser Boston, Boston, MA, 1991.
[39] J.-L. Waldspurger, *Démonstration d'une conjecture de dualité de Howe dans le cas p-adique, $p \neq 2$,* Festschrift in honor of I. I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Part I (Ramat Aviv, 1989), 267–324, Israel Math. Conf. Proc., 2, Weizmann, Jerusalem, 1990.
[40] N. R. Wallach, *Lie Algebra Cohomology and Holomorphic Continuation of Generalized Jacquet Integrals*, Advanced Studies in Math, vol 14, (1988), 123–151.
[41] N. R. Wallach, "Real Reductive Groups I", Academic Press, Boston, 1988.
[42] N. R. Wallach, "Real Reductive Groups II", Academic Press, San Diego, 1992.
[43] H. Yamashita, *On Whittaker vectors for generalized Gelfand-Graev representations of semisimple Lie groups,* J. Math. Kyoto Univ. 26, (1986), no. 2, 263–298.
[44] H. Yamashita, "Cayley transform and generalized Whittaker models for irreducible highest weight modules" in *Nilpotent Orbits, Associated Cycles and Whittaker Models for Highest weight Representations,* Astérisque 273, (2001), 81–137.

Department of Mathematics, National University of Singapore, Block S17, 10 Lower Kent Ridge Road, Singapore 119076
E-mail address: `matrgm@nus.edu.sg`

Department of Mathematics, National University of Singapore, Block S17, 10 Lower Kent Ridge Road, Singapore 119076
E-mail address: `matzhucb@nus.edu.sg`